

## 9.0 REFERENCES

### 9.1 Glossary

**Adult stage:** One of several marked phases or periods in the development and growth of many animals. In vertebrates, the life history stage where the animal is capable of reproducing, as opposed to the juvenile stage.

**Adverse effect:** Any impact that reduces quality and/or quantity of EFH. May include direct or indirect physical, chemical, or biological alterations of the waters or substrate and loss of, or injury to, benthic organisms, prey species and their habitat, and other ecosystem components, if such modifications reduce the quality and or quantity of EFH. Adverse effects to EFH may result from actions occurring within EFH or outside of EFH and may include sites-specific or habitat wide impacts, including individual, cumulative, or synergistic consequences of actions.

**Aggregation:** A group of animals or plants occurring together in a particular location or region.

**Anadromous species:** fish that spawn in fresh or estuarine waters and migrate to ocean waters

**Amphipods:** A small crustacean of the order Amphipoda, such as the beach flea, having a laterally compressed body with no carapace.

**Anaerobic sediment:** Sediment characterized by the absence of free oxygen.

**Anemones:** Any of numerous flowerlike marine coelenterates of the class Anthozoa, having a flexible cylindrical body and tentacles surrounding a central mouth.

**Annual total mortality:** Rate of death expressed as the fraction of a cohort dying over a period compared to the number alive at the beginning of the period ( $\#$  total deaths during year / numbers alive at the beginning of the year). Optimists convert death rates into annual survival rate using the relationship  $S=1-A$ .

**ASPIC (A Surplus Production Model Incorporating Covariates):** A non-equilibrium surplus production model developed by Prager (1995). ASPIC was frequently used by the Overfishing Definition Panel to define  $B_{MSY}$  and  $F_{MSY}$  reference points. The model output was also used to estimate rebuilding timeframes for the Amendment 9 control rules.

**Bay:** An inlet of the sea or other body of water usually smaller than a gulf; a small body of water set off from the main body; e.g. Ipswich Bay in the Gulf of Maine.

**Benthic community:** *Benthic* means the bottom habitat of the ocean, and can mean anything as shallow as a salt marsh or the intertidal zone, to areas of the bottom that are several miles deep in the ocean.

## REFERENCES

### Glossary

*Benthic community* refers to those organisms that live in and on the bottom. (In meaning they live within the substrate; e.g, within the sand or mud found on the bottom. See *Benthic infauna*, below)

**Benthic infauna:** See *Benthic community*, above. Those organisms that live *in* the bottom sediments (sand, mud, gravel, etc.) of the ocean. As opposed to *benthic epifauna*, that live *on* the surface of the bottom sediments.

**Benthivore:** Usually refers to fish that feed on benthic or bottom dwelling organisms.

**Berm:** A narrow ledge typically at the top or bottom of a slope; e.g. a berm paralleling the shoreline caused by wave action on a sloping beach; also an elongated mound or wall of earth.

**Biogenic habitats:** Ocean habitats whose physical structure is created or produced by the animals themselves; e.g, coral reefs.

**Biomass:** The total mass of living matter in a given unit area or the weight of a fish stock or portion thereof. Biomass can be listed for beginning of year (Jan-1), Mid-Year, or mean (average during the entire year). In addition, biomass can be listed by age group (numbers at age \* average weight at age) or summarized by groupings (e.g., age 1<sup>+</sup>, ages 4+ 5, etc). See also spawning stock biomass, exploitable biomass, and mean biomass.

**B<sub>MSY</sub>:** The stock biomass that would produce MSY when fished at a fishing mortality rate equal to F<sub>MSY</sub>. For most stocks, B<sub>MSY</sub> is about 1/2 of the carrying capacity. The proposed overfishing definition control rules call for action when biomass is below 1/4 or 1/2 B<sub>MSY</sub>, depending on the species.

**B<sub>threshold</sub>:** 1) A limit reference point for biomass that defines an unacceptably low biomass i.e., puts a stock at high risk (recruitment failure, depensation, collapse, reduced long term yields, etc). 2) A biomass threshold that the SFA requires for defining when a stock is overfished. A stock is overfished if its biomass is below B<sub>threshold</sub>. A determination of overfished triggers the SFA requirement for a rebuilding plan to achieve B<sub>target</sub> as soon as possible, usually not to exceed 10 years except certain requirements are met. In Amendment 9 control rules, B<sub>threshold</sub> is often defined as either 1/2B<sub>MSY</sub> or 1/4 B<sub>MSY</sub>. B<sub>threshold</sub> is also known as B<sub>minimum</sub>.

**B<sub>target</sub>:** A desirable biomass to maintain fishery stocks. This is usually synonymous with B<sub>MSY</sub> or its proxy.

**Biomass weighted F:** A measure of fishing mortality that is defined as an average of fishing mortality at age weighted by biomass at age for a ranges of ages within the stock (e.g., ages 1<sup>+</sup> biomass weighted F is a weighted average of the mortality for ages 1 and older, age 3<sup>+</sup> biomass weighted is a weighted average for ages 3 and older). Biomass weighted F can also be calculated using catch in weight over mean biomass. See also fully-recruited F.

**Biota:** All the plant and animal life of a particular region.

**Bivalve:** A class of mollusks having a soft body with platelike gills enclosed within two shells hinged together; e.g., clams, mussels.

## REFERENCES

### Glossary

**Bottom roughness:** The inequalities, ridges, or projections on the surface of the seabed that are caused by the presence of bedforms, sedimentary structures, sedimentary particles, excavations, attached and unattached organisms, or other objects; generally small scale features.

**Bottom tending mobile gear:** All fishing gear that operates on or near the ocean bottom that is actively worked in order to capture fish or other marine species. Some examples of bottom tending mobile gear are otter trawls and dredges.

**Bottom tending static gear:** All fishing gear that operates on or near the ocean bottom that is not actively worked; instead, the effectiveness of this gear depends on species moving to the gear which is set in a particular manner by a vessel, and later retrieved. Some examples of bottom tending static gear are gillnets, traps, and pots.

**Boulder reef:** An elongated feature (a chain) of rocks (generally piled boulders) on the seabed.

**Bryozoans:** Phylum aquatic organisms, living for the most part in colonies of interconnected individuals. A few to many millions of these individuals may form one colony. Some bryozoans encrust rocky surfaces, shells, or algae others form lacy or fan-like colonies that in some regions may form an abundant component of limestones. Bryozoan colonies range from millimeters to meters in size, but the individuals that make up the colonies are rarely larger than a millimeter. Colonies may be mistaken for hydroids, corals or seaweed.

**Burrow:** A hole or excavation in the sea floor made by an animal (as a crab, lobster, fish, burrowing anemone) for shelter and habitation.

**Bycatch:** (v.) the capture of nontarget species in directed fisheries which occurs because fishing gear and methods are not selective enough to catch only target species; (n.) fish which are harvested in a fishery but are not sold or kept for personal use, including economic discards and regulatory discards but not fish released alive under a recreational catch and release fishery management program.

**Capacity:** the level of output a fishing fleet is able to produce given specified conditions and constraints. Maximum fishing capacity results when all fishing capital is applied over the maximum amount of available (or permitted) fishing time, assuming that all variable inputs are utilized efficiently.

**Catch:** The sum total of fish killed in a fishery in a given period. Catch is given in either weight or number of fish and may include landings, unreported landings, discards, and incidental deaths.

**Closed Area Model:** A General Algebraic Modeling System (GAMS) model used to evaluate the effectiveness of effort controls used in the Northeast Multispecies Fishery. Using catch data from vessels in the fishery, the model estimates changes in exploitation that may result from changes in DAS, closed areas, and possession limits. These changes in exploitation are then converted to changes in fishing mortality to evaluate proposed measures.

**Coarse sediment:** Sediment generally of the sand and gravel classes; not sediment composed primarily of mud; but the meaning depends on the context, e.g. within the mud class, silt is coarser than clay.

**Commensalism:** See *Mutualism*. An interactive association of two species where one benefits in some way, while the other species is in no way affected by the association.

**Continental shelf waters:** The waters overlying the continental shelf, which extends seaward from the shoreline and deepens gradually to the point where the sea floor begins a slightly steeper descent to the deep ocean floor; the depth of the shelf edge varies, but is approximately 200 meters in many regions.

**Control rule:** A pre-determined method for determining fishing mortality rates based on the relationship of current stock biomass to a biomass target. Amendment 9 overfishing control rules define a target biomass ( $B_{MSY}$  or proxy) as a management objective. The biomass threshold ( $B_{threshold}$  or  $B_{min}$ ) defines a minimum biomass below which a stock is considered overfished.

**Cohort:** see yearclass.

**Crustaceans:** Invertebrates characterized by a hard outer shell and jointed appendages and bodies. They usually live in water and breathe through gills. Higher forms of this class include lobsters, shrimp and crawfish; lower forms include barnacles.

**Days absent:** an estimate by port agents of trip length. This data was collected as part of the NMFS weighout system prior to May 1, 1994.

**Days-at-sea (DAS):** the total days, including steaming time that a boat spends at sea to fish. Amendment 13 categorized DAS for the multispecies fishery into three categories, based on each individual vessel's fishing history during the period fishing year 1996 through 2001. The three categories are: Category A: can be used to target any groundfish stock; Category B: can only be used to target healthy stocks; Category C: cannot be used until some point in the future. Category B DAS are further divided equally into Category B (regular) and Category B (reserve).

**DAS "flip":** A practice in the Multispecies FMP that occurs when a vessel fishing on a Category B (regular) DAS must change ("flip") its DAS to a Category A DAS because it has exceeded a catch limit for a stock of concern.

**Demersal species:** Most often refers to fish that live on or near the ocean bottom. They are often called benthic fish, groundfish, or bottom fish.

**Diatoms:** Small mobile plants (algæ) with silicified (silica, sand, quartz) skeletons. They are among the most abundant phytoplankton in cold waters, and an important part of the food chain.

**Discards:** animals returned to sea after being caught; see Bycatch (n.)

**Dissolved nutrients:** Non-solid nutrients found in a liquid.

**Echinoderms:** A member of the Phylum Echinodermata. Marine animals usually characterized by a five-fold symmetry, and possessing an internal skeleton of calcite plates, and a complex water vascular system. Includes echinoids (sea urchins), crinoids (sea lillies) and asteroids (starfish).

**Ecosystem-based management:** a management approach that takes major ecosystem components and services—both structural and functional—into account, often with a multispecies or habitat perspective

**Egg stage:** One of several marked phases or periods in the development and growth of many animals. The life history stage of an animal that occurs after reproduction and refers to the developing embryo, its food store, and sometimes jelly or albumen, all surrounded by an outer shell or membrane. Occurs before the *larval* or *juvenile stage*.

**Elasmobranch:** Any of numerous fishes of the class Chondrichthyes characterized by a cartilaginous skeleton and placoid scales: sharks; rays; skates.

**Embayment:** A bay or an indentation in a coastline resembling a bay.

**Emergent epifauna:** See *Epifauna*. Animals living upon the bottom that extend a certain distance above the surface.

**Epifauna:** See *Benthic infauna*. *Epifauna* are animals that live on the surface of the substrate, and are often associated with surface structures such as rocks, shells, vegetation, or colonies of other animals.

**Essential Fish Habitat (EFH):** Those waters and substrate necessary to fish for spawning, breeding, feeding, or growth to maturity. The EFH designation for most managed species in this region is based on a legal text definition and geographical area that are described in the Habitat Omnibus Amendment (1998).

**Estuarine area:** The area of an estuary and its margins; an area characterized by environments resulting from the mixing of river and sea water.

**Estuary:** A water passage where the tide meets a river current; especially an arm of the sea at the lower end of a river; characterized by an environment where the mixing of river and seawater causes marked variations in salinity and temperature in a relatively small area.

**Eutrophication:** A set of physical, chemical, and biological changes brought about when excessive nutrients are released into the water.

**Euphotic zone:** The zone in the water column where at least 1% of the incident light at the surface penetrates.

**Exclusive Economic Zone (EEZ):** a zone in which the inner boundary is a line coterminous with the seaward boundary of each of the coastal States and the outer boundary is line 200 miles away and parallel to the inner boundary

**Exempt fisheries:** Any fishery determined by the Regional Director to have less than 5 percent regulated species as a bycatch (by weight) of total catch according to 50 CFR 648.80(a)(7).

**Exploitable biomass:** The biomass of fish in the portion of the population that is vulnerable to fishing.

## REFERENCES

### Glossary

**Exploitation pattern:** Describes the fishing mortality at age as a proportion of fully recruited  $F$  (full vulnerability to the fishery). Ages that are fully vulnerable experience 100% of the fully recruited  $F$  and are termed fully recruited. Ages that are only partially vulnerable experience a fraction of the fully recruited  $F$  and are termed partially recruited. Ages that are not vulnerable to the fishery (including discards) experience no mortality and are considered pre-recruits. Also known as the partial recruitment pattern, partial recruitment vector or fishery selectivity.

**Exploitation rate ( $u$ ):** The fraction of fish in the exploitable population killed during the year by fishing. This is an annual rate compared to  $F$ , which is an instantaneous rate. For example, if a population has 1,000,000 fish large enough to be caught and 550,000 are caught (landed and discarded) then the exploitation rate is 55%.

**Fathom:** A measure of length, containing six feet; the space to which a man can extend his arms; used chiefly in measuring cables, cordage, and the depth of navigable water by soundings.

**Fishing mortality ( $F$ ):** A measurement of the rate of removal of fish from a population caused by fishing. This is usually expressed as an instantaneous rate ( $F$ ) and is the rate at which fish are harvested at any given point in a year. Instantaneous fishing mortality rates can be either fully recruited or biomass weighted. Fishing mortality can also be expressed as an exploitation rate (see exploitation rate) or less commonly, as a conditional rate of fishing mortality ( $m$ , fraction of fish removed during the year if no other competing sources of mortality occurred. Lower case  $m$  should not be confused with upper case  $M$ , the instantaneous rate of natural mortality).

**$F_{0.1}$ :** a conservative fishing mortality rate calculated as the  $F$  associated with 10 percent of the slope at origin of the yield-per-recruit curve.

**$F_{MAX}$ :** a fishing mortality rate that maximizes yield per recruit.  $F_{MAX}$  is less conservative than  $F_{0.1}$ .

**$F_{MSY}$ :** a fishing mortality rate that would produce MSY when the stock biomass is sufficient for producing MSY on a continuing basis.

**$F_{threshold}$ :** 1) The maximum fishing mortality rate allowed on a stock and used to define overfishing for status determination. Amendment 9 frequently uses  $F_{MSY}$  or  $F_{MSY}$  proxy for  $F_{threshold}$ . 2) The maximum fishing mortality rate allowed for a given biomass as defined by a control rule.

**Fishing effort:** the amount of time and fishing power used to harvest fish. Fishing power is a function of gear size, boat size and horsepower.

**Framework adjustments:** adjustments within a range of measures previously specified in a fishery management plan (FMP). A change usually can be made more quickly and easily by a framework adjustment than through an amendment. For plans developed by the New England Council, the procedure requires at least two Council meetings including at least one public hearing and an evaluation of environmental impacts not already analyzed as part of the FMP.

REFERENCES  
Glossary

**Furrow:** A trench in the earth made by a plow; something that resembles the track of a plow, as a marked narrow depression; a groove with raised edges.

**Glacial moraine:** A sedimentary feature deposited from glacial ice; characteristically composed of unsorted clay, sand, and gravel. Moraines typically are hummocky or ridge-shaped and are located along the sides and at the fronts of glaciers.

**Glacial till:** Unsorted sediment (clay, sand, and gravel mixtures) deposited from glacial ice.

**Grain size:** the size of individual sediment particles that form a sediment deposit; particles are separated into size classes (e.g. very fine sand, fine sand, medium sand, among others); the classes are combined into broader categories of mud, sand, and gravel; a sediment deposit can be composed of few to many different grain sizes.

**Growth overfishing:** Fishing at an exploitation rate or at an age at entry that reduces potential yields from a cohort but does not reduce reproductive output (see recruitment overfishing).

**Halocline:** The zone of the ocean in which salinity increases rapidly with depth.

**Habitat complexity:** Describes or measures a habitat in terms of the variability of its characteristics and its functions, which can be biological, geological, or physical in nature. Refers to how complex the physical structure of the habitat is. A bottom habitat with *structure-forming organisms*, along with other three dimensional objects such as boulders, is more complex than a flat, featureless, bottom.

**Highly migratory species:** tuna species, marlin, oceanic sharks, sailfishes, and swordfish

**Hydroids:** Generally, animals of the Phylum Cnidaria, Class Hydrozoa; most hydroids are bush-like polyps growing on the bottom and feed on plankton, they reproduce asexually and sexually.

**Immobile epifaunal species:** See *epifauna*. Animals living on the surface of the bottom substrate that, for the most part, remain in one place.

**Individual Fishing Quota (IFQ):** federal permit under a limited access system to harvest a quantity of fish, expressed by a unit or units representing a percentage of the total allowable catch of a fishery that may be received or held for exclusive use by an individual person or entity

**Juvenile stage:** One of several marked phases or periods in the development and growth of many animals. The life history stage of an animal that comes between the *egg* or *larval stage* and the *adult stage*; juveniles are considered immature in the sense that they are not yet capable of reproducing, yet they differ from the larval stage because they look like smaller versions of the adults.

**Landings:** The portion of the catch that is harvested for personal use or sold.

**Land runoff:** The part of precipitation, snowmelt, or irrigation water that reaches streams (and thence the sea) by flowing over the ground, or the portion of rain or snow that does not percolate into the ground and is discharged into streams instead.

**Larvae stage:** One of several marked phases or periods in the development and growth of many animals. The first stage of development after hatching from the *egg* for many fish and invertebrates. This life stage looks fundamentally different than the juvenile and adult stages, and is incapable of reproduction; it must undergo metamorphosis into the juvenile or adult shape or form.

**Lethrinids:** Fish of the genus *Lethrinus*, commonly called emperors or nor'west snapper, are found mainly in Australia's northern tropical waters. Distinctive features of Lethrinids include thick lips, robust canine teeth at the front of the jaws, molar-like teeth at the side of the jaws and cheeks without scales. Lethrinids are carnivorous bottom-feeding fish with large, strong jaws.

**Limited-access permits:** permits issued to vessels that met certain qualification criteria by a specified date (the "control date").

**Lutjanids:** Fish of the genus of the Lutjanidae: snappers. Marine; rarely estuarine. Some species do enter freshwater for feeding. Tropical and subtropical: Atlantic, Indian and Pacific Oceans.

**Macrobenthos:** See *Benthic community* and *Benthic infauna*. Benthic organisms whose shortest dimension is greater than or equal to 0.5 mm.

**Maturity ogive:** A mathematical model used to describe the proportion mature at age for the entire population.  $A_{50}$  is the age where 50% of the fish are mature.

**Mean biomass:** The average number of fish within an age group alive during a year multiplied by average weight at age of that age group. The average number of fish during the year is a function of starting stock size and mortality rate occurring during the year. Mean biomass can be aggregated over several ages to describe mean biomass for the stock. For example the mean biomass summed for ages 1 and over is the  $1^+$  mean biomass; mean biomass summed across ages 3 and over is  $3^+$  mean biomass.

**Megafaunal species:** The component of the fauna of a region that comprises the larger animals, sometimes defined as those weighing more than 100 pounds.

**Mesh selectivity ogive:** A mathematical model used to describe the selectivity of a mesh size (proportion of fish at a specific length retained by mesh) for the entire population.  $L_{25}$  is the length where 25% of the fish encountered are retained by the mesh.  $L_{50}$  is the length where 50% of the fish encountered are retained by the mesh.

**Meter:** A measure of length, equal to 39.37 English inches, the standard of linear measure in the metric system of weights and measures. It was intended to be, and is very nearly, the ten millionth part of the distance from the equator to the north pole, as ascertained by actual measurement of an arc of a meridian.

**Metric ton:** A unit of weight equal to a thousand kilograms (1kgs = 2.2 lbs.). A metric ton is equivalent to 2,205 lbs. A thousand metric tons is equivalent to 2.2 million lbs.

**Microalgal:** Small microscopic types of algae such as the green algae.

**Microbial:** Microbial means of or relating to microorganisms.

**Minimum spawning stock threshold:** the minimum spawning stock size (or biomass) below which there is a significantly lower chance that the stock will produce enough new fish to sustain itself over the long term.

**Mobile organisms:** organisms that are not confined or attached to one area or place, that can move on their own, are capable of movement, or are moved (often passively) by the action of the physical environment (waves, currents, etc.).

**Molluscs:** Common term for animals of the phylum Mollusca. Includes groups such as the bivalves (mussels, oysters etc.), cephalopods (squid, octopus etc.) and gastropods (abalone, snails). Over 80,000 species in total with fossils back to the Cambrian period.

**Mortality:** see Annual total mortality (A), Exploitation rate (u), Fishing mortality (F), Natural mortality (M), and instantaneous total mortality (Z).

**Motile:** Capable of self-propelled movement. A term that is sometimes used to distinguish between certain types of organisms found in water.

**Multispecies:** the group of species managed under the Northeast Multispecies Fishery Management Plan. This group includes whiting, red hake and ocean pout plus the regulated species (cod, haddock, pollock, yellowtail flounder, winter flounder, witch flounder, American plaice, windowpane flounder, white hake and redfish).

**Mutualism:** See *Commensalism*. A symbiotic interaction between two species in which both derive some benefit.

**Natural disturbance:** A change caused by natural processes; e.g. in the case of the seabed, changes can be caused by the removal or deposition of sediment by currents; such natural processes can be common or rare at a particular site.

**Natural mortality:** A measurement of the rate of death from all causes other than fishing such as predation, disease, starvation, and pollution. Commonly expressed as an instantaneous rate (M). The rate of natural mortality varies from species to species, but is assumed to be  $M=0.2$  for the five critical stocks. The natural mortality rate can also be expressed as a conditional rate (termed  $n$  and not additive with competing sources of mortality such as fishing) or as annual expectation of natural death (termed  $v$  and additive with other annual expectations of death).

**Nearshore area:** The area extending outward an indefinite but usually short distance from shore; an area commonly affected by tides and tidal and storm currents, and shoreline processes.

**Nematodes:** a group of elongated, cylindrical worms belonging to the phylum Nematodea, also called thread-worms or eel-worms. Some non-marine species attack roots or leaves of plants, others are parasites on animals or insects.

REFERENCES  
Glossary

**Nemerteans:** Proboscis worms belonging to the phylum Nemertea, and are soft unsegmented marine worms that have a threadlike proboscis and the ability to stretch and contract.

**Nemipterids:** Fishes of the Family Nemipteridae, the threadfin breams or whiptail breams. Distribution: Tropical and sub-tropical Indo-West Pacific.

**Northeast Shelf Ecosystem:** The Northeast U.S. Shelf Ecosystem has been described as including the area from the Gulf of Maine south to Cape Hatteras, extending from the coast seaward to the edge of the continental shelf, including the slope sea offshore to the Gulf Stream.

**Northwest Atlantic Analysis Area (NAAA):** A spatial area developed for analysis purposes only. The boundaries of this the area are within the 500 fathom line to the east, the coastline to the west, the Hague line to the north, and the North Carolina/ South Carolina border to the south. The area is approximately 83,550 square nautical miles, and is used as the denominator in the EFH analysis to determine the percent of sediment, EFH, and biomass contained in an area, as compared to the total NAAA.

**Nutrient budgets:** An accounting of nutrient inputs to and production by a defined ecosystem (e.g., salt marsh, estuary) versus utilization within and export from the ecosystem.

**Observer:** any person required or authorized to be carried on a vessel for conservation and management purposes by regulations or permits under this Act

**Oligochaetes:** See *Polychaetes*. Oligochaetes are worms in the phylum Annelida having bristles borne singly along the length of the body.

**Open access:** describes a fishery or permit for which there is no qualification criteria to participate. Open-access permits may be issued with restrictions on fishing (for example, the type of gear that may be used or the amount of fish that may be caught).

**Opportunistic species:** Species that colonize disturbed or polluted sediments. These species are often small, grow rapidly, have short life spans, and produce many offspring.

**Optimum Yield (OY):** the amount of fish which A) will provide the greatest overall benefit to the nation, particularly with respect to food production and recreational opportunities, and taking into account the protection of marine ecosystems; B) is prescribed as such on the basis of the maximum sustainable yield from the fishery, as reduced by any relevant economic, social, or ecological factor; and C) in the case of an overfished fishery, provides for rebuilding to a level consistent with producing the maximum sustainable yield in such fishery

**Organic matter:** Material of, relating to, or derived from living organisms.

**Overfished:** A conditioned defined when stock biomass is below minimum biomass threshold and the probability of successful spawning production is low.

REFERENCES  
Glossary

**Overfishing:** A level or rate of fishing mortality that jeopardizes the long-term capacity of a stock or stock complex to produce MSY on a continuing basis.

**Peat bank:** A bank feature composed of partially carbonized, decomposed vegetable tissue formed by partial decomposition of various plants in water; may occur along shorelines.

**Pelagic gear:** Mobile or static fishing gear that is not fixed, and is used within the water column, not on the ocean bottom. Some examples are mid-water trawls and pelagic longlines.

**Phytoplankton:** Microscopic marine plants (mostly algae and diatoms) which are responsible for most of the photosynthetic activity in the oceans.

**Piscivore:** A species feeding preferably on fish.

**Planktivore:** An animal that feeds on plankton.

**Polychaetes:** Polychaetes are segmented worms in the phylum Annelida. Polychaetes (poly-chaetae = many-setae) differ from other annelids in having many setae (small bristles held in tight bundles) on each segment.

**Porosity:** The amount of free space in a volume of a material; e.g. the space that is filled by water between sediment particles in a cubic centimeter of seabed sediment.

**Possession-limit-only permit:** an open-access permit (see above) that restricts the amount of multispecies a vessel may retain (currently 500 pounds of "regulated species").

**Pre-recruits:** Fish in size or age groups that are not vulnerable to the fishery (including discards).

**Prey availability:** The availability or accessibility of prey (food) to a predator. Important for growth and survival.

**Primary production:** The synthesis of organic materials from inorganic substances by photosynthesis.

**Recovery time:** The period of time required for something (e.g. a habitat) to achieve its former state after being disturbed.

**Recruitment:** the amount of fish added to the fishery each year due to growth and/or migration into the fishing area. For example, the number of fish that grow to become vulnerable to fishing gear in one year would be the recruitment to the fishery. "Recruitment" also refers to new year classes entering the population (prior to recruiting to the fishery).

**Recruitment overfishing:** fishing at an exploitation rate that reduces the population biomass to a point where recruitment is substantially reduced.

**Regulated groundfish species:** cod, haddock, pollock, yellowtail flounder, winter flounder, witch flounder, American plaice, windowpane flounder, white hake and redfish. These species are usually targeted with large-mesh net gear.

**Relative exploitation:** an index of exploitation derived by dividing landings by trawl survey biomass. This measure does not provide an absolute magnitude of exploitation but allows for general statements about trends in exploitation.

**Retrospective pattern:** A pattern of systematic over-estimation or underestimation of terminal year estimates of stock size, biomass or fishing mortality compared to that estimate for that same year when it occurs in pre-terminal years.

**Riverine area:** The area of a river and its banks.

**Saurids:** Fish of the family Scomberesocidae, the sauries or needlefishes. Distribution: tropical and temperate waters.

**Scavenging species:** An animal that consumes dead organic material.

**Sea whips:** A coral that forms long flexible structures with few or no branches and is common on Atlantic reefs.

**Sea pens:** An animal related to corals and sea anemones with a featherlike form.

**Sediment:** Material deposited by water, wind, or glaciers.

**Sediment suspension:** The process by which sediments are suspended in water as a result of disturbance.

**Sedentary:** See *Motile* and *Mobile organisms*. Not moving. Organisms that spend the majority of their lives in one place.

**Sedimentary bedforms:** Wave-like structures of sediment characterized by crests and troughs that are formed on the seabed or land surface by the erosion, transport, and deposition of particles by water and wind currents; e.g. ripples, dunes.

**Sedimentary structures:** Structures of sediment formed on the seabed or land surface by the erosion, transport, and deposition of particles by water and wind currents; e.g. ripples, dunes, buildups around boulders, among others.

**Sediment types:** Major combinations of sediment grain sizes that form a sediment deposit, e.g. mud, sand, gravel, sandy gravel, muddy sand, among others.

**Spawning adult stage:** See *adult stage*. Adults that are currently producing or depositing eggs.

**Spawning stock biomass (SSB):** the total weight of fish in a stock that sexually mature, i.e., are old enough to reproduce.

**Species assemblage:** Several species occurring together in a particular location or region

**Species composition:** A term relating the relative abundance of one species to another using a common measurement; the proportion (percentage) of various species in relation to the total on a given area.

**Species diversity:** The number of different species in an area and their relative abundance

**Species richness:** See *Species diversity*. A measurement or expression of the number of species present in an area; the more species present, the higher the degree of species richness.

**Species with vulnerable EFH:** If a species was determined to be “highly” or “moderately” vulnerable to bottom tending gears (otter trawls, scallop dredges, or clam dredges) then it was included in the list of species with vulnerable EFH. Currently there are 23 species and life stages that are considered to have vulnerable EFH for this analysis.

**Status Determination:** A determination of stock status relative to  $B_{\text{threshold}}$  (defines overfished) and  $F_{\text{threshold}}$  (defines overfishing). A determination of either overfished or overfishing triggers a SFA requirement for rebuilding plan (overfished), ending overfishing (overfishing) or both.

**Stock:** A grouping of fish usually based on genetic relationship, geographic distribution and movement patterns. A region may have more than one stock of a species (for example, Gulf of Maine cod and Georges Bank cod). A species, subspecies, geographical grouping, or other category of fish capable of management as a unit.

**Stock assessment:** determining the number (abundance/biomass) and status (life-history characteristics, including age distribution, natural mortality rate, age at maturity, fecundity as a function of age) of individuals in a stock

**Stock of concern:** a regulated groundfish stock that is overfished, or subject to overfishing.

**Structure-forming organisms:** Organisms, such as corals, colonial bryozoans, hydroids, sponges, mussel beds, oyster beds, and seagrass that by their presence create a three-dimensional physical structure on the bottom. See *biogenic habitats*.

**Submerged aquatic vegetation:** Rooted aquatic vegetation, such as seagrasses, that cannot withstand excessive drying and therefore live with their leaves at or below the water surface in shallow areas of estuaries where light can penetrate to the bottom sediments. SAV provides an important habitat for young fish and other aquatic organisms.

**Surficial sediment:** Sediment forming the sea floor or land surface; thickness of the surficial layer may vary.

**Surplus production:** Production of new stock biomass defined by recruitment plus somatic growth minus biomass loss due to natural deaths. The rate of surplus production is directly proportional to stock biomass and its relative distance from the maximum stock size at carrying capacity (K).  $B_{MSY}$  is often defined as the biomass that maximizes surplus production rate.

**Surplus production models:** A family of analytical models used to describe stock dynamics based on catch in weight and CPUE time series (fishery dependent or survey) to construct stock biomass history. These models do not require catch at age information. Model outputs may include stock biomass history, biomass weighted fishing mortality rates,  $MSY$ ,  $F_{MSY}$ ,  $B_{MSY}$ ,  $K$ , (maximum population biomass where stock growth and natural deaths are balanced) and  $r$  (intrinsic rate of increase).

**Survival rate (S):** Rate of survival expressed as the fraction of a cohort surviving the a period compared to number alive at the beginning of the period (# survivors at the end of the year / numbers alive at the beginning of the year). Pessimists convert survival rates into annual total mortality rate using the relationship  $A=1-S$ .

**Survival ratio (R/SSB):** an index of the survivability from egg to age-of-recruitment. Declining ratios suggest that the survival rate from egg to age-of-recruitment is declining.

**TAC:** Total allowable catch. This value is calculated by applying a target fishing mortality rate to exploitable biomass.

**Taxa:** The plural of taxon. Taxon is a named group or organisms of any rank, such as a particular species, family, or class.

**Ten-minute- “squares” of latitude and longitude (TMS):** Are a measure of geographic space. The actual size of a ten-minute-square varies depending on where it is on the surface of the earth, but in general each square is approximately 70-80 square nautical miles in this region. This is the spatial area that EFH designations, biomass data, and some of the effort data have been binned into for analysis purposes in various sections of this document.

**Topography:** The depiction of the shape and elevation of land and sea floor surfaces.

**Total Allowable Catch (TAC):** The amount (in metric tons) of a stock that is permitted to be caught during a fishing year. In the Multispecies FMP, TACs can either be “hard” (fishing ceases when the TAC is caught) or a “target” (the TAC is merely used as an indicator to monitor effectiveness of management measures, but does not trigger a closure of the fishery).

**Total mortality:** The rate of mortality from all sources (fishing, natural, pollution) Total mortality can be expressed as an instantaneous rate (called  $Z$  and equal to  $F + M$ ) or Annual rate (called  $A$  and calculated as the ratio of total deaths in a year divided by number alive at the beginning of the year)

**Trophic guild:** Trophic is defined as the feeding level within a system that an organism occupies; e.g., predator, herbivore. A guild is defined as a group of species that exploit the same class of environmental resources in a similar way. The trophic guild is a utilitarian concept covering both structure and organization that exists between the structural categories of trophic groups and species.

**Turbidity:** Relative water clarity; a measurement of the extent to which light passing through water is reduced due to suspended materials.

**Two-bin (displacement) model:** a model used to estimate the effects of area closures. This model assumes that effort from the closed areas (first bin) is displaced to the open areas (second bin). The total effort in the system is then applied to the landings-per-unit-effort (LPUE) in open areas to obtain a projected catch. The percent reduction in catch is calculated as a net result.

**Vulnerability:** In order to evaluate the potential adverse effects of fishing on EFH, the vulnerability of each species EFH was determined. This analysis defines vulnerability as the likelihood that the functional value of EFH would be adversely affected as a result of fishing with different gear types. A number of criteria were considered in the evaluation of the vulnerability of EFH for each life stage including factors like the function of habitat for shelter, food and/or reproduction.

**Yield-per-recruit (YPR):** the expected yield (weight) of individual fish calculated for a given fishing mortality rate and exploitation pattern and incorporating the growth characteristics and natural mortality.

**Yearclass:** also called cohort. Fish that were spawned in the same year. By convention, the “birth date” is set to January 1st and a fish must experience a summer before turning 1. For example, winter flounder that were spawned in February-April 1997 are all part of the 1997 cohort (or year-class). They would be considered age 0 in 1997, age 1 in 1998, etc. A summer flounder spawned in October 1997 would have its birth date set to the following January 1 and would be considered age 0 in 1998, age 1 in 1999, etc.

**Z:** instantaneous rate of total mortality. The components of Z are additive (i.e.,  $Z = F+M$ )

**Zooplankton:** See *Phytoplankton*. Small, often microscopic animals that drift in currents. They feed on detritus, phytoplankton, and other zooplankton. They are preyed upon by fish, shellfish, whales, and other zooplankton.

REFERENCES  
Glossary

## 9.2 Literature Cited

- Aguilar, R., J. Mas, and X. Pastor. 1995. Impact of Spanish swordfish longline fisheries on the loggerhead sea turtle, *Caretta caretta*, population in the western Mediterranean. U.S. Dep. Commer. NOAA Tech. Memo. NOAA Fisheries-SEFSC-361:1-6.
- Alverson, Dayton L. 1998. Discarding practices and unobserved fishing mortality in marine fisheries: an update. Sea Grant, Washington, DC.
- Balazs, G.H. 1985. Impact of ocean debris on marine turtles: entanglement and ingestion. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-54:387-429.
- Barlow, J., and P. J. Clapham. 1997. A new birth-interval approach to estimating demographic parameters of humpback whales. *Ecology*, 78: 535-546.
- Bass, A.L., S.P. Epperly, J.Braun, D.W. Owens, and R.M. Patterson. 1998. Natal origin and sex ratios of foraging sea turtles in Pamlico-Abermarle Estuarine Complex. U.S. Dep. Commer. NOAA Tech. Memo. NMFS-SEFSC-415.
- Bass, A.L., S.P. Epperly, J.Braun-McNeill. 2004. Multi-year analysis of stock composition of a loggerhead sea turtle (*Caretta caretta*) foraging habitat using maximum likelihood and Bayesian methods. *Conserv. Genetics* 5:783-796.
- Baumgartner, M.F., and B.R. Mate. 2005. Summer and fall habitat of North Atlantic right whales (*Eubalaena glacialis*) inferred from satellite telemetry. *Can. J. Fish. Aquat. Sci.* 62:527-543.
- Berube, M. and A.Aguilar. 1998. A new hybrid between a blue whale, *Balaenoptera musculus*, and a fin whale, *B. physalus*: frequency and implications of hybridization. *Mar. Mamm. Sci.* 14:82-98.
- Best, P.B., J. L., Brownell, R.L. Jr., and Donovan, G.P., (eds.) 2001. Report of the workshop on status and trends of western North Atlantic right whales. *J. Cetacean Res. Manage.* (Special Issue) 2: 61-87.
- Bjorndal, K.A. 1997. Foraging ecology and nutrition of sea turtles. Pp. 199-233 In: Lutz, P.L. and J.A. Muscik, eds. *The Biology of Sea Turtles*. CRC Press, New York. 432 pp.
- Blaylock, R.A., J.W. Hain, L.J. Hansen, D.L. Palka, and G.T. Waring. 1995. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments. NOAA Tech. Memo. NMFS-SEFSC-363. U.S. Department of Commerce, Washington, D.C. 211 pp.
- Bolten, A.B., K.A. Bjorndal, and H.R. Martins. 1994. Life history model for the loggerhead sea turtle (*Caretta caretta*) populations in the Atlantic: Potential impacts of a longline fishery. U.S. Dep. Commer. NOAA Tech. Memo. NOAA Fisheries-SWFSC-201:48-55.
- Boulon, R., Jr., 2000. Trends in sea turtle strandings, U.S. Virgin Islands: 1982 to 1997. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SEFSC-436:261-263.
- Braun, J., and S.P. Epperly. 1996. Aerial surveys for sea turtles in southern Georgia waters, June 1991. *Gulf of Mexico Science*. 1996(1): 39-44.
- Braun-McNeill, J., and S.P. Epperly. 2004. Spatial and temporal distribution of sea turtles in the western North Atlantic and the U.S. Gulf of Mexico from Marine Recreational Fishery Statistics Survey (MRFSS). *Mar. Fish. Rev.* 64(4):50-56.

REFERENCES  
Literature Cited

- Bresette, M.J., R.M. Herren, and D.A. Singewald. 2003. Sea turtle captures at the St. Lucie nuclear power plant: a 25-year synopsis. P. 46. *In*: J.A. Seminoff (compiler). Proceedings of the Twenty-Second Annual Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-503, 308 p.
- Brown, M.B., O.C. Nichols, M.K. Marx, and J.N. Ciano. Surveillance of North Atlantic right whales in Cape Cod Bay and adjacent waters. Final report to the Division of Marine Fisheries, Commonwealth of Massachusetts. 29 pp., September 2002.
- Brown, S.G. 1986. Twentieth-century records of right whales (*Eubalaena glacialis*) in the northeast Atlantic Ocean. *In*: R.L. Brownell Jr., P.B. Best, and J.H. Prescott (eds.) Right whales: Past and Present Status. IWC Special Issue No. 10. p. 121-128.
- Carr, A.R. 1963. Pan specific reproductive convergence in *Lepidochelys kempi*. *Ergebn. Biol.* 26: 298-303.
- Castroviejo, J., J.B. Juste, J.P. Del Val, R. Castelo, and R. Gil. 1994. Diversity and status of sea turtle species in the Gulf of Guinea islands. *Biodiversity and Conservation* 3:828-836.
- Caswell, H., M. Fujiwara, and S. Brault. 1999. Declining survival probability threatens the North Atlantic right whale. *Proc. Nat. Acad. Sci.* 96: 3308-3313.
- Cetacean and Turtle Assessment Program (CeTAP). 1982. Final report of the cetacean and turtle assessment program, University of Rhode Island, to Bureau of Land Management, U.S. Department of the Interior. Ref. No. AA551-CT8-48. 568 pp.
- Chevalier, J., X. Desbois, and M. Girondot. 1999. The reason for the decline of leatherback turtles (*Dermochelys coriacea*) in French Guiana: a hypothesis p.79-88. *In* Miaud, C. and R. Guyétant (eds.), Current Studies in Herpetology, Proceedings of the ninth ordinary general meeting of the Societas Europea Herpetologica, 25-29 August 1998 Le Bourget du Lac, France.
- Clapham, P.J. S. Brault, H. Caswell, M. Fujiwara, S. Kraus, R. Pace, and P. Wade. Report of the working group on survival estimation of North Atlantic right whales. September 27, 2002.
- Cole, T.V., D.L. Hartley, and R.L. Merrick.. 2005. Mortality and Serious Injury Determinations for Large Whale Stocks Along the Eastern Seaboard of the United States, 1999-2003. U.S. Dep. Commer., Northeast Fish. Sci. Cent. Ref. Doc. 05-18. 18p.
- Cole, T., D. Hartley and M. Garron 2006. Mortality and Serious Injury Determinations for Baleen Whale Stocks Along the Eastern Seaboard of the United States, 2000-2004. U.S. Dep. Commer. Northeast Fish. Sci. Cent. Ref. Doc. 06-04. 8p.
- Colvocoresses, J.A. and J.A. Musik. 1983. Species associations and community composition of middle Atlantic bight continental shelf demersal fishes. *U.S. Fisheries Bulletin* 82(2):295-313.
- Crouse, D.T. 1999. The consequences of delayed maturity in a human-dominated world. *American Fisheries Society Symposium*. 23:195-202.
- Crowder, L.B., S.R. Hopkins-Murphy, and A. Royle. 1995. Estimated effect of turtle excluder devices (TEDs) on loggerhead sea turtle strandings with implications for conservation. *Copeia*. 1995:773-779.
- DeAlteris, J., L. Skrobe and C. Lipsky. 1999. The significance of seabed disturbance by mobile fishing gear relative to natural processes: a case study in Narragansett Bay, Rhode Island. *In* L.R. Benaka, editor. Fish Habitat: Essential fish habitat and rehabilitation. American Fisheries Society, Symposium 22, Bethesda, Maryland.

REFERENCES  
Literature Cited

- Department of Fisheries and Oceans, Canada. 1992. Cod and Haddock Separator Trawl. Industry Services and Native Fisheries Report No. 38, November 1992. Scotia-Fundy Region, Fisheries and Oceans, Canada.
- Di Jin and Thunberg, Eric. 2005. An analysis of fishing vessel accidents in fishing area off the northeastern United States. *Safety Science*. 43:523-540.
- Dodd, C.K. 1988. Synopsis of the biological data on the loggerhead sea turtles *Caretta caretta* (Linnaeus 1758). U.S. Fish and Wildlife Service, Biological Report 88 (14).
- Doughty, R.W. 1984. Sea turtles in Texas: A forgotten commerce. *Southwestern Historical Quarterly*. pp. 43-70.
- Dwyer, K.L., C.E. Ryder, and R. Prescott. 2002. Anthropogenic mortality of leatherback sea turtles in Massachusetts waters. Poster presentation for the 2002 Northeast Stranding Network Symposium.
- Eckert, S.A. and J. Lien. 1999. Recommendations for eliminating incidental capture and mortality of leatherback sea turtles, *Dermochelys coriacea*, by commercial fisheries in Trinidad and Tobago. A report to the Wider Caribbean Sea Turtle Conservation Network (WIDECAST). Hubbs-Sea World Research Institute Technical Report No. 2000-310, 7 pp.
- Eckert, S.A., D.W. Nellis, K.L. Eckert, and G.L. Kooyman. 1996. Diving Patterns of Two Leatherback Sea Turtles, (*Demochelys coriacea*) During Interesting Intervals at Sandy Point, St. Croix, U.S. Virgin Islands. *Herpetologica*. Sep. 42(3):381-388.
- Ehrhart, L.M. 1979. A survey of marine turtle nesting at Kennedy Space Center, Cape Canaveral Air Force Station, North Brevard County, Florida, 1-122. Unpublished report to the Division of Marine Fisheries, St. Petersburg, Florida, Florida Department of Natural Resources.
- Engås, A., Jørgensen, T. and West, C. W. 1998. A species-selective trawl for demersal gadoid fisheries. *ICES Journal of Marine Science*, 55: 835-845.
- Epperly, S., L. Avens, L. Garrison, T. Henwood, W. Hoggard, J. Mitchell, J. Nance, J. Poffenberger, C. Sasso, E. Scott-Denton, and C. Yeung. 2002. Analysis of sea turtle bycatch in the commercial shrimp fisheries of southeast U.S. waters and the Gulf of Mexico. U.S. Department of Commerce, NOAA Tech. Memo. NMFS-SEFSC-490, 88pp.
- Epperly, S.P. and W.G. Teas. 2002. Turtle Excluder Devices - Are the escape openings large enough? *Fish. Bull.* 100:466-474.
- Epperly, S.P., J. Braun, and A.J. Chester. 1995a. Aerial surveys for sea turtles in North Carolina inshore waters. *Fishery Bulletin* 93:254-261.
- Epperly, S.P., J. Braun, A.J. Chester, F.A. Cross, J. Merriner, and P.A. Tester. 1995b. Winter distribution of sea turtles in the vicinity of Cape Hatteras and their interactions with the summer flounder trawl fishery. *Bull. Mar. Sci.* 56(2):519-540.
- Epperly, S.P., J. Braun, and A. Veishlow. 1995c. Sea turtles in North Carolina waters. *Cons. Biol.* 9(2): 384-394.
- Ernst, C.H. and R.W. Barbour. 1972. *Turtles of the United States*. Univ. Press of Kentucky, Lexington. 347 pp.
- Farrington, Marriane. Selectivity and Survival of Atlantic Cod and Haddock in the Northwest Atlantic Longline Fishery. NOAA Project 95-NER-141. July 31, 1998.

REFERENCES  
Literature Cited

- Frazer, N.B. and L.M. Ehrhart. 1985. Preliminary growth models for green, *Chelonia mydas*, and loggerhead, *Caretta caretta*, turtles in the wild. *Copeia*. 1985-73-79.
- Frazer, N.B., C.J. Limpus, and J.L. Greene. 1994. Growth and age at maturity for Queensland loggerheads. U.S. Department of Commerce, NOAA Tech. Memo. NMFS-SEFSC-351: 42-45.
- Fritts, T.H. 1982. Plastic bags in the intestinal tracts of leatherback marine turtles. *Herpetological Review* 13(3): 72-73.
- Fujiwara, M. and H. Caswell. 2001. Demography of the endangered North Atlantic right whale. *Nature* 414: 537-541.
- Gabriel, W. 1992. Persistence of demersal fish assemblages between Cape Hatteras and Nova Scotia, northwest Atlantic. *Journal of Northwest Atlantic Fisheries Science* 14:29-46.
- Gilbert, J.R. and N. Guldager. 1998. Status of harbor and gray seal populations in northern New England. Final Report to NMFS, NEFSC, Woods Hole, MA. Coop. Agree. 14-16-009-1557. 13pp.
- Girondot, M., M.H. Godfrey, and R. Philippe. in press. Historical records and trends of leatherbacks in French Guiana and Suriname. *Chelonian Conservation and Biology*.
- Goff, G.P. and J.Lien. 1988. Atlantic leatherback turtle, *Dermochelys coriacea*, in cold water off Newfoundland and Labrador. *Can. Field Nat.*102(1):1-5.
- Graff, D. 1995. Nesting and hunting survey of the turtles of the island of São Tomé. Progress Report July 1995, ECOFAC Componente de São Tomé e Príncipe, 33 pp.
- Hall-Arber, Madeleine, Christopher Dyer, John Poggie, James McNally and Renee Gagne. 2001. Fishing Communities and Fishing Dependency in the Northeast Region of the United States. MARFIN Project Final Report to National Marine Fisheries Service.
- Halliday, R.G., C. G. Cooper, P. Fanning, W. M. Hickey, and P. Gagnon. 1999. Size selection of Atlantic cod, haddock, and pollock (saithe) by otter trawls with square and diamond mesh codends of 130-155 mm mesh size. *Fisheries Research* 41: 255-271.
- Hamilton, P.K., and C.A. Mayo. 1990. Population characteristics of right whales (*Eubalaena glacialis*) observed in Cape Cod and Massachusetts Bays, 1978-1986. *Rep. Int. Whal. Comm.*, Special Issue 12: 203-208.
- Hamilton, P.K., M.K. Marx, and S.D. Kraus. 1998. Scarification analysis of North Atlantic right whales (*Eubalaena glacialis*) as a method of assessing human impacts. Final report to the Northeast Fisheries Science Center, NMFS, Contract No. 4EANF-6-0004.
- Hartley, Marcus. 2004. A review of the economic analyses contained in the DSEIS for Amendment 13. [http://www.nefsc.noaa.gov/groundfish/Hartley\\_review\\_DSEIS.pdf](http://www.nefsc.noaa.gov/groundfish/Hartley_review_DSEIS.pdf).
- Heppell, S.S., L.B. Crowder, D.T. Crouse, S.P. Epperly, and N.B. Frazer. 2003. Population models for Atlantic loggerheads: Past, Present, and Future. *In*: Bolten, A.B. and B.E. Witherington (eds.) *Loggerhead Sea Turtles*. Smithsonian Institution.
- Hilterman, M.L. and E. Goverse. 2004. Annual report of the 2003 leatherback turtle research and monitoring project in Suriname. World Wildlife Fund - Guianas Forests and Environmental Conservation Project (WWF-GFECF) Technical Report of the Netherlands Committee for IUCN (NC-IUCN), Amsterdam, the Netherlands, 21p.
- Hirth, H.F. 1971. Synopsis of biological data on the green sea turtle, *Chelonia mydas*. FAO Fisheries Synopsis No. 85: 1-77.

REFERENCES  
Literature Cited

- Hutchings, J.A., T. D. Bishop, and C. R. McGregor-Shaw. 1999. Spawning behaviour of Atlantic cod, *Gadus morhua*: evidence of mate competition and mate choice in a broadcast spawner. *Canadian Journal of Fisheries and Aquatic Sciences* 56: 97-104.
- ICES 2001. Effects of Different Types of Fisheries on North Sea and Irish Sea Benthic Ecosystems. Report of the ICES Advisory Committee on the Marine Environment 2000. ICES Coop. Res. Rep. No. 241, 27 pp.
- International Council for the Exploration of the Seas [ICES]. 2000. Report of the Working Group on Ecosystem Effects of Fishing Activities. ICES CM 2000/ACE:2.
- International Whaling Commission [IWC]. 2001. Report of the workshop on the comprehensive assessment of right whales: A worldwide comparison. Reports of the International Whaling Commission. Special Issue 2.
- IWC. 1992. Report of the comprehensive assessment special meeting on North Atlantic fin whales. *Rep. Int. Whal. Commn* 42:595-644.
- Johnson, K.A. 2002. A review of national and international literature on the effects of fishing on benthic habitats. NOAA Technical Memorandum NMFS-F/SPO-57. 72 p.
- Katona, S.K., and J.A. Beard. 1990. Population size, migrations, and feeding aggregations of the humpback whale (*Megaptera novaeangliae*) in the Western North Atlantic Ocean. *Rep. Int. Whal. Comm.*, Special Issue 12: 295-306.
- Keinath, J.A., J.A. Musick, and R.A. Byles. 1987. Aspects of the biology of Virginia's sea turtles: 1979-1986. *Virginia J. Sci.* 38(4):329-336.
- Kenney, R.D. 2000. Are right whales starving? Electronic newsletter of the Center for Coastal Studies, posted at [www.coastalstudies.org/entanglementupdate/kenney1.html](http://www.coastalstudies.org/entanglementupdate/kenney1.html) on November 29, 2000. 5pp.
- Kenney, R.D. 2002. North Atlantic, North Pacific, and Southern Hemisphere right whales, pp.806-813 in: W.F. Perrin, B. Wursig, and J.G.M. Thewissen (eds.) *Encyclopedia of Marine Mammals*. Academic Press, CA 2002.
- Kenney, R.D., M.A.M. Hyman, R.E. Owen, G.P. Scott, and H.E. Winn. 1986. Estimation of prey densities required by Western North Atlantic right whales. *Mar. Mamm. Sci.* 2(1): 1-13.
- Kjesbu, O.S., P. Solemdal, P. Bratland, and M. Fonn. 1996. Variation in annual egg production in individual captive Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences* 53: 610-620.
- Knowlton, A. R., J. Sigurjonsson, J.N. Ciano, and S.D. Kraus. 1992. Long-distance movements of North Atlantic right whales (*Eubalaena glacialis*). *Mar. Mamm. Sci.* 8(4): 397-405.
- Knowlton, A.R., S.D. Kraus, and R.D. Kenney. 1994. Reproduction in North Atlantic right whales (*Eubalaena glacialis*). *Can. J. Zool.* 72: 1297-1305.
- Knutsen, G. M. and S. Tisleth. 1985. Growth, development, and feeding success of Atlantic cod larvae (*Gadus morhua*) related to egg size. *Transactions of the American Fisheries Society*, 114:507-511.
- Kraus S. D., Moira W. Brown, Hal Caswell, Christopher W. Clark, Masami Fujiwara, Philip K. Hamilton, Robert D. Kenney, Amy R. Knowlton, Scott Landry, Charles A. Mayo, William A. McLellan, Michael J. Moore, Douglas P. Nowacek, D. Ann Pabst, Andrew J. Read, Rosalind M. Rolland. 2005. North Atlantic Right Whales in Crisis. *Science*, Vol. 309 No. 5734, pp. 561-562.

REFERENCES  
Literature Cited

- Kraus, S.D., M.J. Crone, and A.R. Knowlton 1988. The North Atlantic right whale, in: W.E. Chandler (ed.). The Audubon Wildlife Report 1988/1989. Academic Press, N.Y.
- Kraus, S.D., P.K. Hamilton, R.D. Kenney, A.R. Knowlton, and C.K. Slay. 2001. Reproductive parameters of the North Atlantic right whale. *J. Cetacean Res. Manage.* 2: 231-236.
- Lageux, C.J., C. Campbell, L.H. Herbst, A.R. Knowlton and B. Weigle. 1998. Demography of marine turtles harvested by Miskitu Indians of Atlantic Nicaragua. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SEFSC-412:90.
- Leatherwood, S., and R.R. Reeves. 1983. The Sierra Club handbook of whales and dolphins. Sierra Club Books, San Francisco, California. 302 pp.
- Lewison, R.L., S.A. Freeman, and L.B. Crowder. 2004. Quantifying the effects of fisheries on threatened species: the impact of pelagic longlines on loggerhead and leatherback sea turtles. *Ecology Letters.* 7: 221-231.
- Lutcavage, M.E. and P. Plotkin, B. Witherington, and P.L. Lutz. 1997. Human impacts on sea turtle survival, p.387-409. *In:* P.L. Lutz and J.A. Musick, (eds.), *The Biology of Sea Turtles*, CRC Press, Boca Raton, Florida. 432pp.
- Mahon, R., S.K. Brown, K.C.T. Zwanenburg, D.B. Atkinson, K.R. Buja, L. Claflin, G.D. Howell, M.E. Monaco, R.N. O'Boyle, and M. Sinclair. 1998. Assemblages and biogeography of demersal fishes of the East Coast of North America. *Can. J. Fish. Aquat. Sci.* 55: 1704-1738
- Malik, S., M. W. Brown, S.D. Kraus and B. N. White. 2000. Analysis of mitochondrial DNA diversity within and between North and South Atlantic right whales. *Mar. Mammal Sci.* 16:545-558.
- Marcano, L.A. and J.J. Alio-M. 2000. Incidental capture of sea turtles by the industrial shrimping fleet off northwestern Venezuela. U.S. department of Commerce, NOAA Technical Memorandum NMFS-SEFSC-436:107.
- Márquez, R. 1990. *FAO Species Catalogue, Vol. 11. Sea turtles of the world, an annotated and illustrated catalogue of sea turtle species known to date.* FAO Fisheries Synopsis, 125. 81pp.
- Mate, B.M., S.L. Nieu Kirk, and S.D. Kraus. 1997. Satellite monitored movements of the North Atlantic right whale. *J. Wildl. Manage.* 61:1393-1405.
- Mayo, C. A. and M. K. Marx. 1990. Surface foraging behavior of the North Atlantic right whale., *Eubalaena glacialis*, and associated zooplankton characteristics. *Can. J. Zool.* 68:2214-2220.
- McCay, Bonnie and Marie Cieri. 2000. *Fishing Ports of the Mid-Atlantic.* Report to the Mid-Atlantic Fishery Management Council. Dover, Delaware.
- Meylan, A., B. Schroeder, and A. Mosier. 1995. Sea turtle nesting activity in the state of Florida. *Fla. Mar. Res. Publ.* 52:1-51.
- Milton, S.L., S. Leone-Kabler, A.A. Schulman, and P.L. Lutz. 1994. Effects of Hurricane Andrew on the sea turtle nesting beaches of South Florida. *Bulletin of Marine Science*, 54-3:974-981.
- Mitchell, E. and D.G. Chapman. 1977. Preliminary assessment of stocks of northwest Atlantic sei whales (*Balaenoptera borealis*). *Rep. Int. Whal. Comm. Special Edition* 1:117-120.

REFERENCES  
Literature Cited

- Mizroch, S.A. and A.E. York. 1984. Have pregnancy rates of Southern Hemisphere fin whales, *Balaenoptera physalus*, increased? Rep. Int. Whal. Commn (Spec. Iss. 6):401-410.
- Moore, M.J., A.R. Knowlton, S.D. Krauss, W.A. McLellan, and R.K. Bonde. 2004. Morphometry, Gross Morphology and Available Histopathology in North Atlantic Right Whale (*Eubalaena glacialis*) Mortalities (1970-2002). J. Cetacean Res. Manage. 6(3): 199-214.
- Morgan, L.E. and R. Chuenpagdee. 2003. Shifting Gears: Addressing the collateral impacts of fishing methods in U.S. waters, Pew Science Series on Conservation and the Environment, Washington D.C., Island Press, 41 p.
- Morreale, S.J. and E.A. Standora. 1998. Early life stage ecology of sea turtles in northeastern U.S. waters. U.S. Dep. Commer. NOAA Tech. Mem. NOAA Fisheries-SEFSC-413, 49 pp.
- Morreale, S.J. and E.A. Standora. 1998. Vying for the same resources: potential conflict along migratory corridors. Proceedings of the Seventeenth Annual Sea Turtle Symposium. U.S. Dep. Commer. NOAA Tech Memo. NMFS-SEFSC-415. 294pp.
- Mrosovsky, N. 1981. Plastic jellyfish. Marine Turtle Newsletter 17:5-6.
- Murison, L.D., and D.E. Gaskin. 1989. The distribution of right whales and zooplankton in the Bay of Fundy, Canada. Can. J. Zool. 67:1411-1420.
- Murphy, T.M. and S.R. Hopkins. 1984. Aerial and ground surveys of marine turtle nesting beaches in the southeast region. United States Final Report to NMFS-SEFSC. 73pp.
- Musick, J.A. and C.J. Limpus. 1997. Habitat utilization and migration in juvenile sea turtles. Pp 137-164 In: Lutz, P.L. and J.A. Musick, eds., The Biology of Sea Turtles. CRC Press, New York. 432 pp.
- National Marine Fisheries Service. 2002. Interim Action to Implement Measures to Reduce Overfishing of the Northeast Fishery Complex Under the Northeast Multispecies Fishery Management Plan.
- National Research Council. 2002. Effects of Trawling and Dredging on Seafloor Habitat. National Academy Press. 126 p.
- New England Fishery Management Council (NEFMC). 1994. Amendment 5 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council (NEFMC). 1996. Amendment 7 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council (NEFMC). 1998. Amendment 11 to the Northeast Multispecies Fishery Management Plan, Amendment 9 to the Atlantic Sea Scallop Fishery Management Plan, Amendment 1 to the Monkfish Fishery Management Plan, Amendment 1 to the Atlantic Salmon Fishery Management Plan, Components of the Atlantic Herring Fishery Management Plan, For Essential Fish Habitat.
- New England Fishery Management Council (NEFMC). 20005b. Skate Annual Review.
- New England Fishery Management Council (NEFMC). 2003. Amendment 13 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council (NEFMC). 2004a. Framework Adjustment 40A to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council (NEFMC). 2004b. Framework Adjustment 40B to the Northeast Multispecies Fishery Management Plan.

REFERENCES  
Literature Cited

- New England Fishery Management Council (NEFMC). 2004c. Final Fishery Management Plan for the Northeast Skate Complex.
- New England Fishery Management Council (NEFMC). 2005a. Framework Adjustment 41 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council (NEFMC). 2006. Framework Adjustment 43 to the Northeast Multispecies Fishery Management Plan.
- NMFS 1994. State and federal fishery interactions with sea turtles in the Mid-Atlantic area. NOAA/NMFS, Silver Spring, MD, 13pp.
- NMFS 2003. Techniques for Making Weak Links and Marking Buoy Lines. How to Comply with the Atlantic Large Whale Take Reduction Plan. Brochure October, 2003.
- NMFS and U.S. Fish and Wildlife Service (USFWS). 1998. Recovery Plan for U.S. Pacific Populations of the Green Turtle (*Chelonia mydas*). National Marine Fisheries Service, Silver Spring, Maryland.
- NMFS and U.S. Fish and Wildlife Service (USFWS). 1991a. Recovery plan for U.S. population of loggerhead turtle. National Marine Fisheries Service, Washington, D.C. 64 pp.
- NMFS and U.S. Fish and Wildlife Service (USFWS). 1991b. Recovery plan for U.S. population of Atlantic green turtle. National Marine Fisheries Service, Washington, D.C. 58 pp.
- NMFS and U.S. Fish and Wildlife Service (USFWS). 1992. Recovery plan for leatherback turtles in the U.S. Caribbean, Atlantic, and Gulf of Mexico. National Marine Fisheries Service, Washington, D.C. 65 pp.
- NMFS and U.S. Fish and Wildlife Service (USFWS). 1995. Status reviews for sea turtles listed under the Endangered Species Act of 1973. National Marine Fisheries Service, Silver Spring, Maryland. 139 pp.
- NMFS and U.S. Fish and Wildlife Service (USFWS). 1998. Recovery Plan for the U.S. Pacific Population of the Leatherback Turtle (*Dermochelys coriacea*). National Marine Fisheries Service, Silver Spring, Maryland.
- NMFS and USFWS. 1992. Recovery plan for leatherback turtles in the U.S. Caribbean, Atlantic, and Gulf of Mexico. National Marine Fisheries Service, Washington, D.C. 65 pp.
- NMFS Southeast Fisheries Science Center. 2001. Stock assessments of loggerheads and leatherback sea turtles and an assessment of the impact of the pelagic longline fishery on the loggerhead and leatherback sea turtles of the Western North Atlantic. U.S. Department of Commerce, National Marine Fisheries Service, Miami, FL, SEFSC Contribution PRD-00/01-08; Parts I-III and Appendices I-IV. NOAA Tech. Memo NMFS-SEFSC-455, 343 pp.
- NMFS. 2002. Endangered Species Act Section 7 Consultation on Shrimp Trawling in the Southeastern United States, under the Sea Turtle Conservation Regulations and as Managed by the Fishery Management Plans for Shrimp in the South Atlantic and Gulf of Mexico. December 2.
- NMFS. 2004. Endangered Species Act Section 7 Reinitiated Consultation on the Continued Authorization of the Atlantic Pelagic Longline Fishery under the Fishery Management Plan for Atlantic Tunas, Swordfish, and Sharks (HMS FMP). Biological Opinion, June 1.
- NMFS. 1991. Final recovery plan for the North Atlantic right whale (*Eubalaena glacialis*). Prepared by the Right Whale Recovery Team for the National Marine Fisheries Service, Silver Spring, Maryland. 86 pp.

REFERENCES  
Literature Cited

- NMFS. 1998a. Draft recovery plans for the fin whale (*Balaenoptera physalus*) and sei whale (*Balaenoptera borealis*). Prepared by R.R. Reeves, G.K. Silber, and P.M. Payne for the National Marine Fisheries Service, Silver Spring, Maryland. July 1998.
- NMFS. 1998b. Recovery plan for the blue whale (*Balaenoptera musculus*). Prepared by Reeves, R.R., P.J. Clapham, and R.L. Brownell, Jr. for the National Marine Fisheries Service, Silver Spring, Maryland. Mitchell, E. 1974. Present status of the northwest Atlantic fin and other whale stocks. Pages 108-169 in W. E. Schevill (ed) *The Whale Problem: A status report*. Harvard University Press. Cambridge, Massachusetts, 419pp.
- Northeast Fisheries Science Center (NEFSC). 2002a. Assessment of 20 Northeast Groundfish Stocks through 2001: a report of the Groundfish Assessment Review Meeting (GARM), Northeast fisheries Science Center, Woods Hole, MA, October 8-11, 2001.
- Northeast Fisheries Science Center (NEFSC). 2003. Report of the 37th Northeast Regional Stock Assessment Workshop (35th SAW): Consensus Summary of Assessments. NEFSC Reference Document 03-016).
- Northeast Fisheries Science Center (NEFSC). 2005. 40th Northeast Regional Stock Assessment Workshop (40th SAW) Assessment Report. ( NEFSC Reference Document 05-04).
- Northeast Fisheries Science Center (NEFSC). 2005a. Assessment of 19 Northeast Region Groundfish Stocks through 2004: Groundfish Assessment Review Meeting (2005 GARM; GARM II). NEFSC Reference Document 05-13.
- Northeast Fisheries Science Center (NEFSC). 2005b. NEFSC bycatch estimation methodology: allocation, precision, and accuracy, by P.J. Rago, S.E. Wigley, and M.J. Fogarty. NEFSC Reference Document 05-09.
- Northeast Region Essential Fish Habitat Steering Committee (NREFHSC). 2002. Workshop on the Effects of Fishing Gear on Marine Habitats Off the Northeastern United States, October 23-25, 2001, Boston, Massachusetts. Northeast Fish Sci Cent Ref Doc 02-01; 86.
- Norton, John. September 25, 2003. President, Cozy Harbor Seafood, Maine. Personal communication.
- O'Brien, Loretta. 1999. Factors influencing the rate of sexual maturity and the effect on spawning stock for Georges Bank and Gulf of Maine Atlantic cod (*Gadus morhua*) stocks. *Journal of Northwest Atlantic Fisheries Science* 25: 179-203.
- Overholtz, W.J. and A.V. Tyler. 1985. Long-term responses of the demersal fish assemblages of Georges Bank. *U.S. Fisheries Bulletin* 83(4):507-520.
- Palka, D. 2000. Abundance and distribution of sea turtles estimated from data collected during cetacean surveys. *In*: Bjorndal, K.A. and A.B. Bolten. *Proceedings of a workshop on assessing abundance and trends for in-water sea turtle populations*. U.S. Dep. Commer. NOAA Tech. Mem. NMFS-SEFSC-445, 83pp.
- Payne, P.M., D.N. Wiley, S.B. Young, S. Pittman, P.J. Clapham, and J.W. Jossi. 1990. Recent fluctuations in the abundance of baleen whales in the southern Gulf of Maine in relation to changes in selected prey. *Fish. Bull.* 88 (4): 687-696
- Perry, S.L., D.P. DeMaster, and G.K. Silber. 1999. The great whales: History and status of six species listed as endangered under the U.S. Endangered Species Act of 1973. *Mar. Fish. Rev. Special Edition.* 61(1): 59-74.

REFERENCES  
Literature Cited

- Perry, S.L., D.P. DeMaster, and G.K. Silber. 1999. The Sperm Whale In: The great whales: History and status of six species listed as endangered under the U.S. Endangered Species Act of 1973. Mar. Fish. Rev. Special Edition. 61(1): 59-74.
- Pike, B. 2003. The North Atlantic right whale catalog: an update on mortality, reproduction, and population status, in: North Atlantic Right Whale Consortium Annual Meeting Abstracts. New Bedford Whaling Museum, November 4-5. New England Aquarium Compilers.
- Pinhorn, A. T. and R. G. Halliday. 2001. The regulation of exploitation pattern in North Atlantic groundfish stocks. Fisheries Research 53: 25-37.
- Pratt, S. 1973. Benthic fauna. Pp. 5-1 to 5-70 in: Coastal and offshore environmental inventory, Cape Hatteras to Nantucket Shoals. University of Rhode Island, Marine Publication Series No. 2. Kingston, RI.
- Prescott, R.L. 1988. Leatherbacks in Cape Cod Bay, Massachusetts, 1977-1987, p 83-84 In: B.A. Schroeder (comp.), Proceedings of the Eighth Annual Workshop on Sea Turtle Conservation and Biology. NOAA Technical Memorandum NMFS-SEFC-214.
- Pritchard, P.C.H. 1982. Nesting of the leatherback turtle, *Dermochelys coriacea*, in Pacific, Mexico, with a new estimate of the world population status. Copeia 1982:741-747.
- Pritchard, P.C.H. 1997. Evolution, phylogeny and current status. Pp. 1-28 In: The Biology of Sea Turtles. Lutz, P., and J.A. Musick, eds. CRC Press, New York. 432 pp.
- Rankin-Baransky, K., C.J. Williams, A.L. Bass, B.W. Bowen, and J.R. Spotila. 2001. Origin of loggerhead turtles stranded in the northeastern United States as determined by mitochondrial DNA analysis. Journal of Herpetology, v. 35, no. 4, pp 638-646.
- Raymond, John D. and Manomet Center for Conservation Sciences. 2004. A Collaborative Program to Test the Use of a Cod/haddock Separator Panel in Trawl Nets. Report to the NOAA/NMFS CRPI.
- Rebel, T.P. 1974. Sea turtles and the turtle industry of the West Indies, Florida and the Gulf of Mexico. Univ. Miami Press, Coral Gables, Florida.
- Reid, R.N., L.M. Cargnelli, S.J. Griesbach, D.B. Packer, D.L. Johnson, C.A. Zetline, W.W. Morse and P.L. Berrien. 1999. Essential fish habitat source document: (series) [Species] life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-124 through 152. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA.
- Robbins, J., and D. Mattila. 1999. Monitoring entanglement scars on the caudal peduncle of Gulf of Maine humpback whales. Report to the National Marine Fisheries Service. Order No. 40EANF800288. 15 pp.
- Ross, J.P. 1979. Green turtle, *Chelonia mydas*, Background paper, summary of the status of sea turtles. Report to WWF/IUCN. 4pp.
- Schaeff, C.M., Kraus, S.D., Brown, M.W., Perkins, J.S., Payne, R., and White, B.N. 1997. Comparison of genetic variability of North and South Atlantic right whales (*Eubalaena*), using DNA fingerprinting. Can. J. Zool. 75:1073-1080.
- Schevill, W.E., W.A. Watkins, and K.E. Moore. 1986. Status of *Eubalaena glacialis* off Cape Cod. Rep. Int. Whal. Comm., Special Issue 10: 79-82.
- Schultz, J.P. 1975. Sea turtles nesting in Surinam. Zoologische Verhandelingen (Leiden), Number 143: 172 pp.

REFERENCES  
Literature Cited

- Seipt, I., P.J. Clapham, C.A. Mayo, and M.P. Hawvermale. 1990. Population characteristics of individually identified fin whales, *Balaenoptera physalus*, in Massachusetts Bay. *Fish. Bull.* 88:271-278.
- Seminoff, J.A. 2004. *Chelonia mydas*. In: IUCN 2004. 2004 IUCN Red List of Threatened Species. Downloaded on October 12, 2005 from [www.redlist.org](http://www.redlist.org).
- Sherman, K.J., N.A. Jaworski, T.J. Smayda (eds). 1996. The Northeast Shelf Ecosystem – Assessment, Sustainability, and Management. Blackwell Science, Inc. Cambridge, MA.
- Shoop, C.R. and R.D. Kenney. 1992. Seasonal distributions and abundance of loggerhead and leatherback sea turtles in waters of the northeastern United States. *Herpetol. Monogr.* 6: 43-67.
- Skrobe, L. G., Beutel, D. L., Castor, K., Valliere, A., Gibson, M., Lazar, N., Borden, D., Lee, L. 2003. Southern New England Yellowtail Flounder Size Selectivity for 6.0" diamond, 6.5" square, 6.5" diamond, and 7.0" square shaped mesh codends. Technical Report 1-03, Rhode Island Gear Conservation Engineering Working Group, University of Rhode Island Sea Grant.
- Spotila, J.R., A.E. Dunham, A.J. Leslie, A.C. Steyermark, P.T. Plotkin, and F.V. Paladino. 1996. Worldwide Population Decline of *Demochelys coriacea*: Are Leatherback Turtles Going Extinct? *Chelonian Conservation and Biology* 2(2): 209-222.
- Steinback, S. and B. Gentner. 2001. Marine Angler Expenditures in the Northeast Region, 1998. NOAA Technical Memorandum NMFS-F-SPO-47. NMFS Office of Science and Technology, Fisheries Statistics and Economics Division. Silver Spring, MD.
- Steinback, S., and E. Thunberg. In-Press. Northeast region commercial fishing input-output model. NOAA Technical Memorandum NMFS-NE-?.
- Steinback, S.R. 2004. Using ready-made regional input-output models to estimate backward linkage effects of exogenous output shocks. *The Review of Regional Studies*, 34(1):57-71.
- Stephens, S.H. and J. Alvarado-Bremer. 2003. Preliminary information on the effective population size of the Kemp's ridley (*Lepidochelys kempii*) sea turtle. In: Seminoff, J.A., compiler. Proceedings of the Twenty-Second Annual Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-503, 308p.
- Stevenson, D., Chiarella, L., Stephan, D., Reid, R., Wilhelm, K., McCarthy, J. and Pentony, M. 1994 Characterization of the Fishing Practices and Marine Benthic Ecosystems of the Northeast U.S. Shelf, and an Evaluation of the Potential Effects of Fishing on Essential Fish Habitat, NOAA Technical Memorandum NMFS-NE-181.
- Swingle, W.M., S.G. Barco, T.D. Pitchford, W.A. McLellan, and D.A. Pabst. 1993. Appearance of juvenile humpback whales feeding in the nearshore waters of Virginia. *Mar. Mamm. Sci.* 9: 309-315.
- Theroux, R.B. and M.D. Grosslein. 1987. Benthic fauna. Pp. 283-195 in: R.H. Backus (ed.), Georges Bank. MIT Press, Cambridge, MA.
- Theroux, R.B. and R.L. Wigley. 1998. Quantitative composition and distribution of the macrobenthic invertebrate fauna of the continental shelf ecosystems of the northeastern United States. NOAA Technical Report NMFS 140. U.S. Dept. of Commerce, Seattle, WA.

REFERENCES  
Literature Cited

- Thrush, S.F., J.E. Hewitt, V.J. Cummings, P.K. Dayton, M. Cryer, S.J. Turner, G.A. Funnell, R.G. Budd, C.J. Milcurn, and M.R. Wilkinson. 1998. Disturbance of the marine benthic habitat by commercial fishing: impacts at the scale of the fishery. *Ecological Applications* 8(3):866-879.
- Trippel, E. A. and M. J. Morgan. 1994. Age-specific paternal influences on reproductive success of Atlantic cod (*Gadus morhua* L.) of the Grand Banks, Newfoundland. *ICES Mar. Sci. Symp.* 198: 414-422.
- Tschernij, Vesa and Rene Holst. 1999. Evidence of factors at vessel-level affecting codend selectivity in Baltic cod demersal trawl fishery. *ICES CM* 1999/R:02.
- Tuck, I.D., S.J. Hall, M.R. Robertson, E. Armstrong, and D.J. Basford. 1998. Effects of physical trawling disturbance in a previously unfished sheltered Scottish sea loch. *Marine Ecology Progress Series* 162:227-242.
- Turtle Expert Working Group (TEWG). 1998. An assessment of the Kemp's ridley (*Lepidochelys kempii*) and loggerhead (*Caretta caretta*) sea turtle populations in the Western North Atlantic. NOAA Technical Memorandum NMFS-SEFSC-409. 96 pp.
- Turtle Expert Working Group (TEWG). 2000. Assessment update for the Kemp's ridley and loggerhead sea turtle populations in the western North Atlantic. U.S. Dep. Commer. NOAA Tech. Mem. NMFS-SEFSC-444, 115 pp.
- USFWS and NMFS. 2003. Notice of Petition Finding (Fed Register) September 15, 2003.
- USFWS and NMFS. 1992. Recovery plan for the Kemp's ridley sea turtle (*Lepidochelys kempii*). NMFS, St. Petersburg, Florida.
- Valentine, P.C. and R.G. Lough. 1991. The sea floor environment and the fishery of eastern Georges bank. Dept. of Interior, U.S. Geological Survey, Open File Report 91-439.
- Valentine, P.C., E.W. Strom, R.G. Lough, and C.L. Brown. 1993. Maps showing the sedimentary environment of eastern Georges bank. U.S. Geological Survey, Miscellaneous Investigations Series, Map I-2279-B, scale 1:250,000.
- Vallin, L. and A. Nissling. 2000. Maternal effects on egg size and egg buoyancy of Baltic cod, *Gadus morhua*: Implications for stock structure effects on recruitment. *Fisheries Research* 49: 21-37.
- Waring, G.T. Elizabeth Josephson, Carol P. Fairfield, and Katherine Maze-Foley, Editors, *with contributions from (listed alphabetically)*: Dana Beldon, Timothy V.N. Cole, Lance P. Garrison, Keith D. Mullin, Christopher Orphanides, Richard M. Pace, Debra L. Palka, Marjorie C. Rossman, and Fredrick W. Wenzel U.S. Atlantic and Gulf of Mexico marine mammal stock assessments - 2005. NOAA Technical Memorandum NMFS-NE-194
- Waring, G.T., D.L. Palka, P.J. Clapham, S. Swartz, M. Rossman, T. Cole, L.J. Hansen, K.D. Bisack, K. Mullin, R.S. Wells, D.K. Odell, and N.B. Barros. 1999. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments - 1999. NOAA Technical Memorandum NMFS-NE-153.
- Waring, G.T., J.M. Quintal, C. P. Fairfield (eds). 2002. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments - 2002. NOAA Technical Memorandum NMFS-NE-169. 318 p.
- Waring, G.T., R.M. Pace, J.M. Quintal, C. P. Fairfield, K. Maze-Foley (eds). 2004. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments - 2003 . NOAA Technical Memorandum NMFS-NE-182. 287 p.

REFERENCES  
Literature Cited

- Watkins, W.A., and W.E. Schevill. 1982. Observations of right whales (*Eubalaena glacialis*) in Cape Cod waters. *Fish. Bull.* 80(4): 875-880.
- Watkins, W.A., K.E. Moore, J. Sigurjonsson, D. Wartzok, and G. Notarbartolo di Sciara. 1984. Fin whale (*Balaenoptera physalus*) tracked by radio in the Irminger Sea. *Rit Fiskideildar* 8(1): 1-14.
- Watling, L. 1998. Benthic fauna of soft substrates in the Gulf of Maine. Pp. 20-29 in: *Effects of fishing gear on the sea floor of New England*, E.M. Dorsey and J. Pederson (eds.). MIT Sea Grant Pub. 98-4.
- Weisbrod, A.V., D. Shea, M.J. Moore, and J.J. Stegeman. 2000. Organochlorine exposure and bioaccumulation in the endangered Northwest Atlantic right whale (*Eubalaena glacialis*) population. *Environmental Toxicology and Chemistry*, 19(3):654-666.
- Winn, H.E., C.A. Price, and P.W. Sorensen. 1986. The distributional biology of the right whale (*Eubalaena glacialis*) in the western North Atlantic. *Reports of the International Whaling Commission, Special Issue No. 10*:129-138.
- Witzell, W.N. 2002. Immature Atlantic loggerhead turtles (*Caretta caretta*): suggested changes to the life history model. *Herpetological Review* 33(4): 266-269.
- Wynne, K. and M. Schwartz. 1999. *Guide to marine mammals and turtles of the U.S. Atlantic and Gulf of Mexico*. Rhode Island Sea Grant, Narragansett. 115pp.
- Zug, G. R. and J.F. Parham. 1996. Age and growth in leatherback turtles, *Dermochelys coriacea*: a skeletochronological analysis. *Chelonian Conservation and Biology*. 2(2): 244-249.
- Zurita, J.C., R. Herrera, A. Arenas, M.E. Torres, C. Calderon, L. Gomez, J.C. Alvarado, and R. Villavicencio. 2003. Nesting loggerhead and green sea turtles in Quintana Roo, Mexico. Pp. 125-127. In: J.A. Seminoff (compiler). *Proceedings of the Twenty-Second Annual Symposium on Sea Turtle Biology and Conservation*. NOAA Tech. Memo. NMFS-SEFSC-503, 308 p.

## REFERENCES

Index

### 9.3 Index

- Affected Human Environment, 156, 215, 513  
Biological Impacts, 228, 358, 385, 410, 439, 465, 494, 532  
Bycatch, 11, 276, 279, 349, 366, 393, 422, 447, 478, 495, 544, 603, 604  
Cod  
GB, 50, 67, 76, 203, 293, 348, 349, 515, 517  
GOM, 50, 51, 76, 347, 349, 466, 504, 515  
Cod: GOM, 515  
Cumulative Effects, 6, 10, 512, 513, 524, 525, 526, 527, 528, 529, 553, 554, 572, 599  
Days@at@sea (DAS), 221, 514, 515  
Days-at-sea (DAS), 3, 33, 42, 50, 58, 59, 60, 61, 63, 75, 76, 77, 92, 96, 126, 128, 159, 160, 171, 221, 279, 280, 282, 286, 287, 294, 340, 349, 350, 383, 408, 437, 463, 492, 502, 514, 515, 517, 524, 547, 603, 604  
Category A, 3, 57, 58, 61, 76, 77, 92, 286, 287, 517, 524, 604  
Category B, 3, 50, 58, 59, 60, 61, 63, 76, 77, 92, 96, 126, 128, 221, 279, 286, 287, 515, 517, 604  
Category C, 3, 76, 604  
Economic impacts, 6, 156, 220, 304, 327  
Essential Fish Habitat, 6, 7, 33, 75, 95, 99, 126, 128, 129, 140, 159, 280, 282, 284, 289, 513, 520, 522, 530, 533, 551, 552, 553, 554, 555, 601, 605, 610, 613, 614, 615, 621  
Exclusive Economic Zone, 33, 37, 154, 513, 514, 605  
Habitat impacts, 296  
Habitat Impacts, 350, 367, 394, 423, 448, 479, 495, 535  
Haddock  
GB, 76, 77, 203, 347  
GOM, 76, 347  
Halibut, 141, 289  
Incidental Catch TACs, 50, 58, 60, 63, 77, 279, 515  
Magnuson-Stevens Fishery Conservation and Management Act, 3, 34, 37, 42, 76, 156, 276, 327, 340, 513, 517, 519, 524, 547, 549, 553, 554, 572  
Marine mammals, 276, 513, 518, 519, 581, 625  
Ocean pout, 76, 348  
Plaice, American, 48, 50, 60, 76, 96, 132, 140, 141, 285, 289, 609, 612  
Pollock, 48, 60, 141, 289  
Protected species, 143, 513, 518, 526  
Redfish, 48, 60, 76, 141, 289  
Social Impact Analysis, 35, 215, 221  
Social impacts, 9, 327, 357, 408, 437, 462, 491  
Special Access Program, 34  
Closed Area I Hook Gear Haddock SAP, 77, 96, 133  
Closed Area II Haddock SAP, 50, 96, 133, 279  
White hake, 76, 348  
Windowpane flounder, 76, 348  
Winter flounder  
SNE/MA, 76, 203, 348, 521  
Yellowtail flounder  
CC/GOM, 50, 51, 61, 76, 348, 515, 517  
SNE/MA, 60, 61, 76, 348, 516

REFERENCES  
Index

*Intentionally Blank*