

8.0 References

8.1 Literature Cited

- Abernathy, A. (ed). 1989. Description of the Mid-Atlantic environment. U.S. Dept. of the Interior, Minerals Management Service, Herndon, VA. 167 p. plus appendices.
- Able, K.W. and A.M. Muzeni. 2002. An evaluation of the impact of mobile fishing gear on Tilefish (*Lopholatilus chamaeleonticeps*) habitat: review of archived video images from submersibles. Final Report to the Mid-Atlantic Fisheries Management Council. Rutgers University, Institute of Marine and Coastal Science Marine Field Station. Tuckerton, NJ. 28p.
- Able, K.W. and M.P. Fahay. 1998. *The First Year of Life in Estuarine Fishes in the Middle Atlantic Bight*. Rutgers University Press, New Brunswick, NJ.
- Agler, B.A., R.L., Schooley, S.E. Frohock, S.K. Katona, and I.E. Seipt. 1993. Reproduction of photographically identified fin whales, *Balaenoptera physalus*, from the Gulf of Maine. *J. Mamm.* 74:577-587.
- Aguilar, R., J. Mas, and X. Pastor. 1995. Impact of the Spanish swordfish longline fisheries on the loggerhead sea turtle, *Caretta caretta*, population in the western Mediterranean. U.S. Dep. Commer. NOAA Tech Memo. NMFS-SEFSC-361:1-6.
- Aguirre International. 1996. An Appraisal of the Social and Cultural Aspects of the Multispecies Groundfish Fishery in New England and the Mid-Atlantic Regions. Report submitted to the National Oceanic and Atmospheric Administration. Contract Number 50-DGNF-5-00008.
- Almeida, F., L. Arlen, P. Auster, J. Cross, J. Lindholm, J. Link, D. Packer, A. Paulson, R. Reid, and P. Valentine. 2000. The effects of marine protected areas on fish and benthic fauna: the Georges Bank closed area II example. Poster presented at American Fisheries Society 130th Annual Meeting, St. Louis, MO, August 20-24, 2000.
- Alverson, Dayton L. 1998. Discarding practices and unobserved fishing mortality in marine fisheries: an update. Sea Grant, Washington, DC.
- Armstrong, Mike, Margaret Hunter, Josef Idoine, Clare McBane, Dave McCarron, Daniel Schick, and Amy Schick. 2000. Assessment Report for Gulf of Maine Northern Shrimp. Atlantic States Marine Fisheries Commission, Northern Shrimp Technical Committee.
- Anderson, John T. 2001. Classification of marine habitats using submersible and acoustic seabed techniques. In: *Spatial Processes and Management of Marine Populations*. Kruese, Bez, Booth, Dorn, Hills, Lipcius, Pelletier, Roy, Smith, and Witherell (eds). University of Alaska Sea Grant, AK-SG-01-02, Fairbanks, AK, p. 377-394.
- Atkinson, D.B., G.A. Rose, E.F. Murphy, and C.A. Bishop. 1997. Distribution changes and abundance of northern cod (*Gadus morhua*), 1981-1993. *Canadian Journal of Fisheries and Aquatic Science*. 54 (Suppl. 1):123-128.
- Auster, P.J., K. Joy and P.C. Valentine. 2001. Fish species and community distributions as

- proxies for seafloor habitat distributions: the Stellwagen Bank National Marine Sanctuary example (Northwest Atlantic, Gulf of Maine). *Environmental Biology of Fishes* 60:331-346.
- Auster, P.J. and R.W. Langton. 1999. The effects of fishing on fish habitat. In L.R. Benaka, editor. *Fish Habitat: Essential fish habitat and rehabilitation*. American Fisheries Society, Symposium 22, Bethesda, Maryland.
- Auster, P.J. 1998a. A conceptual model of the impacts of fishing gear on the integrity of fish habitats. *Conservation Biology* 12(6): 1198-1203.
- Auster, P.J., C. Michalopoulos, P.C. Valentine, and R.J. Malatesta. 1998b. Delineating and monitoring habitat management units in a temperate deep-water marine protected area. Pages 169-185 in N.W.P. Munro and J.H.M. Willison, editors. *Linking Protected Areas with Working Landscapes, Conserving Biodiversity*. Science and Management of Protected Areas Association, Wolfville, Nova Scotia.
- Auster, P.J., R.J. Malatesta, and C.L.S. Donaldson. 1997. Distributional responses to small-scale variability by early juvenile silver hake, *Merluccius bilinearis*. *Environ. Biol. Fishes* 50: 195-200.
- Auster, P.J., R.J. Malatesta, R.W. Langton, L. Watling, P.C. Valentine, C.L.S. Donaldson, E.W. Langton, A.N. Shepard, and I.G. Babb. 1996. The impacts of mobile fishing gear on seafloor habitats in the Gulf of Maine (northwest Atlantic): implications for conservation of fish populations. *Reviews in Fisheries Science* 4(2):185-202.
- Auster, P.J., R.J. Malatesta, and S.C. LaRosa. 1995. Patterns of microhabitat utilization by mobile megafauna on the southern New England (USA) continental shelf and slope. *Marine Ecology Progress Series*. 127:77-85.
- Auster, P.J., R.J. Malatesta, S.C. LaRosa, R.A. Cooper, and L.L. Stewart. 1991. Microhabitat utilization by the megafaunal assemblage at a low relief outer continental shelf site - Middle Atlantic Bight, USA. *Journal of Northwest Atlantic Fisheries Science*. 11:59-69.
- Auster, P.J., J. Lindholm and P.C. Valentine. In prep. Primary and secondary habitats of juvenile Acadian redfish (*Sebastes faciatius*): patterns of differential survival or elements of a redfish pump? To be submitted to *Environmental Biology of Fishes*.
- Auster, P.J., J. Lindholm, S. Schaub, G. Funnel, L.S. Kaufman, and P.C. Valentine. In press. Use of sand wave habitats by silver hake *Merluccius bilinearis* (Mitchill). *J. Fish Biology*.
- Azarovitz, T.R. 1981. A brief historical review of the Woods Hole Laboratory trawl survey time series, p. 62-67. In: Dobuleday, W.G., and D. Rivard (eds.), *Bottom Trawl Surveys*. Canadian Special Publication in Fisheries and Aquatic Science. 58:273 p.
- Backus, R.H. and D. Bourne (eds.). 1987. *Georges Bank*. Cambridge, MA: MIT Press.
- Ball, B., B. Munday, and I. Tuck. 2000. Effects of otter trawling on the benthos and environment in muddy sediments. Pp. 69-82 in M.J. Kaiser and S.J. de Groot. *The Effects of Fishing on Non-target Species and Habitats*. Blackwell Science.
- Barnette, M.C. 2001. A review of the fishing gear utilized within the Southeast Region and their potential impacts on essential fish habitat. NOAA Technical Memorandum (NMFS-SEFSC-449).

- Barlow, J., and P. J. Clapham. 1997. A new birth-interval approach to estimating demographic parameters of humpback whales. *Ecology*, 78: 535-546. Swingle, W.M., S.G. Barco, T.D. Pitchford, W.A. McLellan, and D.A. Pabst. 1993. Appearance of juvenile humpback whales feeding in the nearshore waters of Virginia. *Mar. Mamm. Sci.* 9: 309-315.
- Barrow, C.J. 1997. *Environmental and Social Impact Assessment: An Introduction*. London: Arnold/Hodder Headline.
- Baum, E. 1997. *Maine Atlantic Salmon, A National Treasure*. Atlantic Salmon Unlimited, Hermon, Maine. 224 pp.
- Beardsley, R.C., B. Butman, W.R. Geyer, and P. Smith. 1996. Physical oceanography of the Gulf of Maine: an update. Pp. 39-52 in: Wallace, G.T. and E.F. Braasch (eds). *Proceedings of the Gulf of Maine Ecosystem Dynamics Scientific Symposium and Workshop*. Regional Association for Research on the Gulf of Maine (RARGOM) Report 97-1.
- Bergman, M.J.N., and J.W. Van Santbrink 2000. Fishing mortality of populations of megafauna in sandy sediments. Pp. 49-68 in M.J. Kaiser and S. J. de Groot. *The effects of fishing on non-target species and habitats*. Blackwell Science.
- Berube, M. and A.Aguilar. 1998. A new hybrid between a blue whale, *Balaenoptera musculus*, and a fin whale, *B. physalus*: frequency and implications of hybridization. *Mar. Mamm. Sci.* 14:82-98.
- Bjorndal, K.A. 1997. Foraging ecology and nutrition of sea turtles. Pp. 199-233 In: Lutz, P.L. and J.A. Musick, eds., *The Biology of Sea Turtles*. CRC Press, New York. 432 pp.
- Black, K.P. and G.D. Parry. 1999. Entrainment, dispersal, and settlement of scallop dredge sediment plumes: field measurements and numerical modelling. *Canadian Journal of Fisheries and Aquatic Science* 56:2271-2281.
- Black, K.P. and G.D. Parry. 1994. Sediment transport rates and sediment disturbance due to scallop dredging in Port Phillip Bay. *Memoirs of the Queensland Museum* 36(2):327-341.
- Blaylock, R.A., J.W. Hain, L.J. Hansen, D.L. Palka, and G.T. Waring. 1995. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments. NOAA Tech. Memo. NMFS-SEFSC-363. U.S. Department of Commerce, Washington, D.C. 211 pp.
- Bockstael, N.E. 1976. Analysis of investment behavior and price determination: analytical input for the formation of policy in fisheries. Ph.D. dissertation. University of Rhode Island, Kingston, RI.
- Boesch, D.F. 1979. Benthic ecological studies: macrobenthos. Chapter 6 in: *Middle Atlantic outer continental shelf environmental studies*. Conducted by Virginia Institute of Marine Studies under contract AA550-CT6062 with the Bureau of Land Management. 301 p.
- Bolten, A.B., K.A. Bjorndal, and H.R. Martins. 1994. Life history model for the loggerhead sea turtle (*Caretta caretta*) populations in the Atlantic: Potential impacts of a longline fishery. US Dept. Commer. NOAA Tech. Memo. NMFS-SEFSC-201:48-55.
- Boulva, J. and I.A. McLaren. 1979. Biology of the harbor seal, *Phoca vitulina*, in eastern Canada. *Bull. Fish. Res. Bd. Can.* 200:1-24.

- Bradshaw, C., L.O. Veale, and A.R. Brand. 2002. The role of scallop-dredge disturbance in long-term changes in Irish Sea benthic communities: a re-analysis of an historical dataset. *Journal of Sea Research* 47:161-184.
- Bradshaw, C., L.O. Veale, A.S. Hill, and A.R. Brand. 2001. The effect of scallop dredging on Irish Sea benthos: experiments using a closed area. *Hydrobiologia* 465:129-138.
- Bradshaw, C., L.O. Veale, A.S. Hill, and A.R. Brand. 2000. The effects of scallop dredging on gravelly seabed communities. Pp. 83-104 in M.J. Kaiser and S.J. de Groot. *The Effects of Fishing on Non-target Species and Habitats*. Blackwell Science.
- Brooks, D.A. 1996. Physical oceanography of the shelf and slope seas from Cape Hatteras to Georges Bank: A brief overview. Pp. 47-75 in: Sherman, K., N.A. Jaworski, and T.J. Smayda (eds). *The Northeast Shelf Ecosystem – Assessment, Sustainability, and Management*. Blackwell Science, Cambridge, MA. 564 p.
- Brown, B. 1993. A classification system of marine and estuarine habitats in Maine: An ecosystem approach to habitats. Part I: Benthic Habitats. Maine Natural Areas Program, Dept. of Economic and Community Development, Augusta, ME. 51 p. + 1 appendix.
- Brylinsky, M., J. Gibson, and D.C. Gordon, Jr. 1994. Impacts of flounder trawls on the intertidal habitat and community of the Minas Basin, Bay of Fundy. *Canadian Journal of Fisheries and Aquatic Sciences* 51:650-661.
- Burdge, R.J. 1998. *A Conceptual Approach to Social Impact Assessment (Revised Edition)*. Madison, WI: Social Ecology Press.
- Burgess, G.H. 2002. Spiny Dogfishes. Family Squalidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Butcher, T., J. Matthews, J. Glaister, and G. Hamer. 1981. Study suggests scallop dredges causing few problems in Jervis Bay. *Australian Fisheries* 40(9):9-12.
- Butman, V., M. Noble and J. Moody 1982. Observations of near-bottom currents at the shelf break near Wilmington Canyon in the Mid-Atlantic outer continental shelf area: results of 1978-1979 field seasons, U.S. Geol. Survey Final Report to U.S. Bureau of Land Management: 3-1-3-58.
- Caddy, J.F. 1973. Underwater observations on tracks of dredges and trawls and some effects of dredging on a scallop ground. *Journal of the Fisheries Research Board of Canada* 30:173-180.
- Caddy, J.F. 1968. Underwater observations on scallop (*Placopecten magellanicus*) behavior and drag efficiency. *Journal of the Fisheries Research Board of Canada* 25(10):2123-2141.
- Cahoon, L.B. 1999. The role of benthic microalgae in neritic ecosystems. *Oceanography and Marine Biology* 37:47-86.
- Canadian Department of Fisheries and Ocean. 1993. Seabed disturbances from fishing activities. Unpublished Report. Canadian Department of Fisheries and Oceans. Scotia-Fundy Region. Industry Services and Native Fisheries Branch. 4 p.
- Cargnelli, L.M., S.J. Griesbach, C. McBride, C.A. Zetlin, and W.W. Morse. 1999a. Essential

- fish habitat source document: longfin inshore squid, *Loligo pealeii*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-146. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 27 p.
- Cargnelli, L.M., S.J. Griesbach, D.B. Packer, and E. Weissberger. 1999b. Essential fish habitat source document: ocean quahog, *Artica islandica*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-148. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 12 p.
- Cargnelli, L.M., S.J. Griesbach, D.B. Packer, and E. Weissberger. 1999c. Essential fish habitat source document: Atlantic surfclam, *Spisula solidissima*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-148. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 13 p.
- Cargnelli, L.M., S.J. Griesbach, D.B. Packer, P.L. Berrien, D.L. Johnson and W.W. Morse. 1999d. Essential fish habitat source document: pollock, *Pollachius virens*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-131. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 30 p.
- Cargnelli, L.M., S.J. Griesbach, D.B. Packer, P.L. Berrien, W.W. Morse and D.L. Johnson. 1999e. Essential fish habitat source document: witch flounder, *Glyptocephalus cynoglossus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-139. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 29 p.
- Cargnelli, L.M., S.J. Griesbach, P.L. Berrien, W.W. Morse, and D.L. Johnson. 1999f. Essential fish habitat source document: haddock, *Melanogrammus aeglefinus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-128. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 31 p.
- Cargnelli, L.M., S.J. Griesbach, and W.W. Morse. 1999g. Essential fish habitat source document: Atlantic halibut, *Hippoglossus hippoglossus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-125. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 17 p.
- Carr, H.A. and H. Milliken. 1998. Conservation engineering: options to minimize fishing's impacts to the sea floor. In: *Effects of Fishing Gear on the Sea Floor of New England* (E.L. Dorsey and J. Pederson eds). Conservation Law Foundation, Boston, Massachusetts.
- Carr, H.A., Pol, M., and Ribas, Luis. 2001. Groundfish Trawlnets desinged to reduce the bycatch of cod. Massachusetts Division of Marine Fisheries.
- Carroll, M.T. 1998. An assessment of the Atlantic bluefin tuna market: Implications for management. Unpublished master's thesis, University of Rhode Island.
- Caruso, J.H. 2002. Goosfishes or Monkfishes. Family Lophiidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Caswell, H., M. Fujiwara, and S. Brault. 1999. Declining survival probability threatens the North Atlantic right whale. *Proc. Nat. Acad. Sci.* 96: 3308-3313.
- Cetacean and Turtle Assessment Program (CeTAP). 1982. Final report or the cetacean and turtle assessment program, University of Rhode Island, to Bureau of Land Management, U.S. Department of the

Interior. Ref. No. AA551-CT8-48. 568 pp.

- Chang, S., P.L. Berrien, D.L. Johnson, and C.A. Zetlin. 1999a. Essential fish habitat source document: offshore hake, *Merluccius albidus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-130. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 24 p.
- Chang, S., P.L. Berrien, D.L. Johnson, and W.W. Morse. 1999b. Essential fish habitat source document: windowpane, *Scopthalmus aquosus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-137. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 32 p.
- Chang, S., W.W. Morse, and P.L. Berrien. 1999c. Essential fish habitat source document: white hake, *Urophycis tenuis*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-136. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 23 p.
- Clapham, P.J. (Ed.) 1999. Predicting right whale distribution, Report of the workshop held on October 1 and 2, 1998, in Woods Hole, Massachusetts. Northeast Fisheries Science Center Reference Document 99-11. 44 pp.
- Clark, C.W. 1995. Application of U.S. Navy underwater hydrophone arrays for scientific research on whales. Rep. Int. Whal. Comm. 45: 210-212.
- Clark, S.H., S.X. Cadrin, D.F. Schick, P.J. Diodati, M.P. Armstrong, and D. McCarron. 2000. Journal of Northwest Atlantic Fisheries Science 27: 193-226.
- Clarke, R. 1954. Open boat whaling in the Azores: the history and present methods of a relic history. Discovery Rep. 26:281-354.
- Coastal Enterprises, Inc. (Moore and Sheehan). May 1998. Perspectives from Five Ports: Impacts of Amendments 5 and 7 to the Northeast Multispecies Fishery Management Plan. Prepared for the Maine Department of Marine Resources.
- Collie, J.S., S.J. Hall, M.J. Kaiser, and I.R. Poiner. 2000. A quantitative analysis of fishing impacts on shelf-sea benthos. Journal of Animal Ecology 69:785-798.
- Collie, J.S., G.A. Escanaero, and P.C. Valentine. 1997. Effects of bottom fishing on the benthic megafauna of Georges Bank. Marine Ecology Progress Series 155:159-172.
- Colvocoresses, J.A. and J.A. Musik. 1983. Species associations and community composition of middle Atlantic bight continental shelf demersal fishes. U.S. Fisheries Bulletin 82(2):295-313.
- Cook, S.K. 1988. Physical oceanography of the middle Atlantic bight. Pp. 1-50 in: A.L. Pacheco (ed). Characterization of the middle Atlantic water management unit of the northeast regional action plan. NOAA Technical Memorandum NMFS-F/NEC-56. Woods Hole, MA. 322 p.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. U.S. Fish and Wildlife Service Publication FWS/OBS-79/31. Washington, DC. 103 p.
- Creaser, E.P., Jr., D.A. Clifford, M.J. Hogan and D.B. Sampson. 1983. A commercial sampling

- program for sandworms, *Nereis virens* Sars, and bloodworms, *Glycera dibranchiata* Ehrens, harvested along the Maine Tidal Coast. NOAA Tech. Rep. NMFS SSRF-767, 56.
- Crouse, D.T. 1999. The consequences of delayed maturity in a human-dominated world. American Fisheries Society Symposium. 23:195-202.
- Currie, D.R. and G.D. Parry. 1999. Impacts and efficiency of scallop dredging on different soft substrates. Canadian Journal of Fisheries and Aquatic Science 56:539-550.
- Currie, D.R. and G.D. Parry. 1996. Effects of scallop dredging on a soft sediment community: a large-scale experimental study. Marine Ecology Progress Series 134:131-150.
- DeAlteris, J., L. Skrobe and C. Lipsky. 1999. The significance of seabed disturbance by mobile fishing gear relative to natural processes: a case study in Narragansett Bay, Rhode Island. In L.R. Benaka, editor. Fish Habitat: Essential fish habitat and rehabilitation. American Fisheries Society, Symposium 22, Bethesda, Maryland.
- DeAlteris, J. 1998. Unpublished Manuscript. Training Manual: Fisheries Science and Technology, prepared for the NOAA Corps Officer Program. University of Rhode Island, Department of Fisheries. Kingston, RI. 34 p.
- DeAlteris, J. and Grogan, C. 1997. An analysis of harvesting gear size selectivity for eight demersal groundfish species in the Northwest Atlantic Ocean. Fisheries Technical Report No. 1, University of Rhode Island.
- DeAlteris, J. and Grogan, C. 1997. An analysis of yield and spawning stock biomass per recruit for eight demersal groundfish species in the northwest Atlantic Ocean. Fisheries Technical Report No. 2, University of Rhode Island.
- Department of Fisheries and Oceans (DFO) Canada. 1992. Cod and haddock separator trawl. Project Summary No. 38, Scotia-Fundy Region, Halifax, Nova Scotia. November 1992.
- Doeringer, Peter, Philip Moss and David Terkla (1986) *The New England Fishing Economy: Jobs, Income, and Kinship*, Amherst: University of Massachusetts Press.
- Dorsey, E.M. 1998. Geological overview of the sea floor of New England. Pp. 8-14 in: Effects of fishing gear on the sea floor of New England, E.M. Dorsey and J. Pederson (eds.). MIT Sea Grant Pub. 98-4.
- DuPaul, W. D., J. E. Kirkley and D. B. Rudders. 1999. Scallop gear selectivity and scallop biology: a mismatch in resource management. Abst. National Shellfisheries Association Annual Meeting, Halifax, Nova Scotia.
- Drabsch, S.L., J.E. Tanner, and S.D. Connell. 2001. Limited infaunal response to experimental trawling in previously untrawled areas. ICES Journal of Marine Science 58:1261-1271.
- Eckert, S.A., D.W. Nellis, K.L. Eckert, and G.L. Kooyman. 1996. Diving Patterns of Two Leatherback Sea Turtles, (*Demochelys coriacea*) During Interesting Intervals at Sandy Point, St. Croix, U.S. Virgin Islands. Herpetologica. Sep. 42(3):381-388.
- Ehrhart, L.M. 1979. A survey of marine turtle nesting at Kennedy Space Center, Cape Canaveral Air Force Station, North Brevard County, Florida, 1-122. Unpublished report to the Div of Mar Fish. St. Pete.,

FL, Flor. Dept. of Nat. Res.

- Eleftheriou, A. and M.R. Robertson. 1992. The effects of experimental scallop dredging on the fauna and physical environment of a shallow sandy community. *Netherlands Journal of Sea Research* 30:289-299.
- Engas, A., T. Jorgensen, and C. W. West. 1998. A species-selective trawl for demersal gadoid fisheries. *ICES Journal of Marine Science* 55: 835-845.
- Engel, J. and R. Kvitek 1998. Effects of otter trawling on a benthic community in Monterey Bay National Marine Sanctuary. *Conservation Biology* 12(6):1204-1214.
- Eno, N.C., D.S. MacDonald, J.A.M. Kinnear, S.C. Amos, C.J. Chapman, R.A. Clark, F.P.D. Bunker, and C. Munro. 2001. Effects of crustacean traps on benthic fauna. *ICES Journal of Marine Science* 58:11-20.
- Epperly, S.P., J. Braun, A.J. Chester, F.A. Cross, J. Merriner, and P.A. Teater. 1995. Winter distribution of sea turtles in the vicinity of Cape Hatteras and their interactions with the summer flounder trawl fishery. *Bull. Mar. Sci.* 56(2):519-540.
- Ernst, C.H. and R.W. Barbour. 1972. *Turtles of the United States*. Univ. Press of Kentucky, Lexington. 347 pp.
- Everhart, W.H. and W.D. Youngs. 1981. *Principles of Fishery Science*, Second Edition. Cornell University Press, Ithaca, NY, 349 pp.
- Fahay, M.P., P.L. Berrien, D.L. Johnson, and W.W. Morse. 1999. Essential fish habitat source document: Atlantic cod, *Gadus morhua*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-124. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 41 p.
- Fonseca, M.S., G.W. Thayer, A.J. Chester and C. Foltz. 1984. Impact of scallop harvesting on eelgrass (*Zostera marina*) meadows: implications for management. *North American Journal of Fisheries Management* 4:286-293.
- Fraser, S., V. Gotceitas, and J. A. Brown. 1996. Interactions between age-classes of Atlantic cod and their distribution among bottom substrates. *Canadian Journal of Fisheries and Aquatic Science* 53:305-314.
- Freese, L., P.J. Auster, J. Heifetz, and B.L. Wing. 1999. Effects of trawling on seafloor habitat and associated invertebrate taxa in the Gulf of Alaska. *Marine Ecology Progress Series* 182:119-126.
- Fretwell, S. and H. Lucas. 1970. On the territorial behavior and other factors influencing habitat distribution in birds. *Acta Biotheoretica* 19: 16-36.
- Frid, C.L.J., R.A. Clark and J.A. Hall. 1999. Long-term changes in the benthos on a heavily fished ground of the NE coast of England. *Marine Ecology Progress Series* 188:13-29.
- Gabriel, W. 1992. Persistence of demersal fish assemblages between Cape Hatteras and Nova Scotia, northwest Atlantic. *Journal of Northwest Atlantic Fisheries Science* 14:29-46.
- Garrison, L.P. 2000. Spatial and dietary overlap in the Georges Bank groundfish community. *Canadian Journal of Fisheries and Aquatic Science*, 57: 1679-1691.

- Garrison, Lance P. 2001. Spatial patterns in species composition in northeast United States continental shelf fish community during 1966-1999. In: *Spatial Processes and Management of Marine Populations*. Kruese, Bez, Booth, Dorn, Hills, Lipcius, Pelletier, Roy, Smith, and Witherell (eds). University of Alaska Sea Grant, AK-SG-01-02, Fairbanks, AK, p. 513-537.
- Gatewood, John B., and Bonnie J. McCay (1988) "Job Satisfaction and the Culture of Fishing: A Comparison of Six New Jersey Fisheries," *Maritime Anthropological Studies* 1(2):103-128.
- _____ (1990) "Comparison of Job Satisfaction in Six New Jersey Fisheries: Implications for Management," *Human Organization* 49(1): 14-25.
- Georgianna, D. and A. Cass. 1998. The costs of hook fishing for groundfish in the Northeastern United States. University of Massachusetts Dartmouth.
- Georgianna, D. and J. Dirlam. 2000. The effect of reduced supply on fish processing in New England. Proceedings of the Tenth Biennial Conference of the International Institute of Fisheries Economics and Trade, July 10-14, 2000. Corvallis, Oregon.
- Georgianna, D., J. Dirlam, and R. Townsend. 1993. The Groundfish and Scallop Processing Sectors in New England. Report Prepared for NMFS.
- Geraci, J.R., D.M. Anderson, R.J. Timperi, D.J. St. Aubin, G.A. Early, J.H. Prescott, and C.A. Mayo. 1990. Humpback whales (*Megaptera novaengliae*) fatally poisoned by dinoflagellate toxin. *Can. J. Fish. And Aquat. Sci.* 46(11):1895-1898.
- Gerstner, C.L. 1998. Use of substratum ripples for flow refugia by Atlantic cod, *Gadus morhua*. *Environmental Biology of Fishes*, 51(4).
- Gerstner, C.L. and P.W. Webb. 1998. The station-holding performance of the plaice *Pleuronectes platessa* on artificial substratum ripples. *Canadian Journal of Zoology*, 76(2).
- Gibbs, P.J., A.J. Collins, and L.C. Collett. 1980. Effect of otter prawn trawling on the macrobenthos of a sandy substratum in a New South Wales estuary. *Australian Journal of Marine and Freshwater Research* 31:509-516.
- Gilbert, J.R. and N. Guldager. 1998. Status of harbor and gray seal populations in northern New England. Final Report to NMFS, NEFSC, Woods Hole, MA. Coop. Agree. 14-16-009-1557. 13pp.
- Gilkinson, K., M. Paulin, S. Hurley, and P. Schwinghamer. 1998. Impacts of trawl door scouring on infaunal bivalves: results of a physical trawl door model/dense sand interaction. *Journal of Experimental marine Biology and Ecology* 224:291-312.
- Glenn, Robert, M. Hunter, J. Idoine, C. McBane, D. McCarron, M. Lewis. 2001. Assessment Report for Gulf of Maine Northern Shrimp. Atlantic State Marine Fisheries Commission, Northern Shrimp Technical Committee.
- Godcharles, M.F. 1971. A study of the effects of a commercial hydraulic clam dredge on benthic

communities in estuarine areas. State of Florida Department of Natural Resources Marine Resources Laboratory Technical Series No. 64.

- Goff, G.P. and J.Lien. 1988. Atlantic leatherback turtle, *Dermochelys coriacea*, in cold water off Newfoundland and Labrador. *Can. Field Nat.* 102(1):1-5.
- Gomez, E.D., A.C. Alcala, and H.T. Yap. 1987. Other fishing methods destructive to coral. In *Human Impacts on Coral Reefs: Facts and Recommendations*. Antenne Museum, French Polynesia. 65-75.
- Graham, N. and R. Kynoch. 2001. Square mesh panels in demersal trawls: some data on haddock selectivity in relation to mesh size and position. *Fisheries Research* 49(2001): 207-218.
- Guthrie, J.F. and C.W. Lewis. 1982. The clam-kicking fishery of North Carolina. *Marine Fisheries Review* 44(1):16-21.
- Hain, J. H. W. 1975. The international regulation of whaling. *Marine Affairs J.* 3: 28-48.
- Hain, J.H.W., M.J. Ratnaswamy, R.D. Kenney, and H.E. Winn. 1992. The fin whale, *Balaenoptera physalus*, in waters of the northeastern United States continental shelf. *Rep. Int. Whal. Comm.* 42: 653-669.
- Hall, S.J. and M. Harding. 1997. Physical disturbance and marine benthic communities: the effects of mechanical harvesting of cockles on non-target benthic infauna. *Journal of Applied Ecology* 34:497-517.
- Hall, S.J., D.J. Basford, and M.R. Robertson. 1990. The impact of hydraulic dredging for razor clams *Ensis* sp. On an infaunal community. *Netherlands Journal of Sea Research* 27(1):119-125.
- Hall, S.J., M.R. Robertson, D.J. Basford, and S.D. Heaney. 1993. The possible effects of fishing disturbance in the northern North Sea: an analysis of spatial patterns in community structure around a wreck. *Netherlands Journal of Sea Research* 31(2):201-208.
- Hall-Arber, M. and D. McCarron. unpublished draft 8 Sept. 2001. Amendment 1 to the Interstate Fishery Management Plan for Northern Shrimp (social and economic sections only). Atlantic States Marine Fisheries Commission.
- Hall-Arber, Madeleine, Christopher Dyer, John Poggie, James McNally and Renee Gagne. 2001. *Fishing Communities and Fishing Dependency in the Northeast Region of the United States*. MARFIN Project Final Report to National Marine Fisheries Service.
- Halliday, R.G., C. G. Cooper, P. Fanning, W. M. Hickey, and P. Gagnon. 1999. Size selection of Atlantic cod, haddock, and pollock (saithe) by otter trawls with square and diamond mesh codends of 130-155 mm mesh size. *Fisheries Research* 41: 255-271.
- Hall-Spencer, J.M. and P.G. Moore. 2000a. Impact of scallop dredging on maerl grounds. In: *Effects of Fishing on Non-Target Species and Habitats: Biological, Conservation, and Socio-economic Issues* (M.J. Kaiser and S.J. de Groot (eds.)). Blackwell Science Ltd., Oxford, England.
- Hamilton, P.K., and C.A. Mayo. 1990. Population characteristics of right whales (*Eubalaena glacialis*) observed in Cape Cod and Massachusetts Bays, 1978-1986. *Rep. Int. Whal. Comm., Special Issue* 12: 203-208.

- Hamilton, P.K., M.K. Marx, and S.D. Kraus. 1998. Scarification analysis of North Atlantic right whales (*Eubalaena glacialis*) as a method of assessing human impacts. Final report to the Northeast Fisheries Science Center, NMFS, Contract No. 4EANF-6-0004.
- Hansson, M., M. Lindegarth, D. Valentinsson and M. Ulmestrand. 2000. Effects of shrimp-trawling on abundance of benthic macrofauna in Gullmarsfjorden, Sweden. *Marine Ecology Progress Series* 198:191-201.
- Harwood, J. 2002. Mass Die-Offs. *In*: W.F. Perrin, B. Würsig, J.G.M. Thewissen (eds.) *Encyclopedia of Marine Mammals*. Academic Press. p. 724-726.
- Hayes, M.L. 1983. Active fish capture methods, in *Fisheries Techniques*. Nielson, L.A. and D.L. Johnson, eds. American Fisheries Society, Bethesda, Maryland.
- Health Care for All (Hams, Herold, & Miller). November 1996. Health Survey of the Fishing Population in Massachusetts, Final Report. Funded by Caritas Christi Health Care System.
- Health Care for All (Hams, Herold, Marra, & Miller). 1999. Health Survey of the Fishing Population in Maine, Final Report. Funded by Caritas Christi Health Care System.
- Hecker, B. 2001. *In*: S. Azimi (ed.), Priority ocean areas for protection in the Mid-Atlantic. Natural Resources Defense Council, Washington, DC. 32-36.
- Hecker, B. 1990. Variation in megafaunal assemblages on the continental margin south of New England. *Deep-Sea Research* 37(1):37-57.
- Hecker, B. and G. Blechschmidt. 1979. Epifauna of the northeastern U.S. continental margin. Appendix A *in*: Hecker, B., G. Blechschmidt, and P. Gibson (eds.), Epifaunal zonation and community structure in three mid- and North Atlantic canyons. Final Report for the Canyon Assessment Study in the Mid- and North Atlantic Areas of the U.S. Outer Continental Shelf. U.S. Dept. of the Interior, Bureau of Land Management, Washington, DC., January 11, 1980.
- Hecker, B., D.T. Logan, F.E. Gandarillas, and P.R. Gibson. 1983. Megafaunal assemblages in canyon and slope habitats. Vol. III: Chapter I. Canyon and Slope Processes Study. Final report prepared for U.S. Dept. of the Interior, Minerals Management Service, Washington, D.C.
- Hicks, R., S. Steinback, A. Gautam, and E. Thunberg. 1999. Volume II: The Economic Value of New England and Mid-Atlantic Sportfishing in 1994. NOAA Technical Memorandum (NMFS-F/SPO-38). NMFS Office of Science and Technology – Fisheries Statistics and Economics Division. Silver Spring, MD.
- Hildebrand, H. 1963. Hallazgo del are de anidacion de la tortuga “lora” *Lepidochelys kempii* (Garman), en la costa occidental del Golfo de Mexico (Rept. Chel.). *Ciencia Mex.*, 22(4):105-112.
- Howell, P.T., D.R. Molnar, and R.B. Harris. 1999. Juvenile winter flounder distribution by habitat type. *Estuaries* 22(4):1090-1095.
- Hubert, W.A. 1983. Passive capture techniques, in *Fisheries Techniques*. Nielson, L.A. and D.L. Johnson, eds. American Fisheries Society, Bethesda, Maryland.
- Hutchings, J.A., T. D. Bishop, and C. R. McGregor-Shaw. 1999. Spawning behaviour of Atlantic

- cod, *Gadus morhua*: evidence of mate competition and mate choice in a broadcast spawner. Canadian Journal of Fisheries and Aquatic Sciences 56: 97-104.
- Inches, Sue. January 2001. Maine Seafood Harvesting and Processing: A Developmental Strategy. Maine Department of Marine Resources.
- IWC. 1971. Report of the Special Meeting on Sperm Whale Biology and Stock Assessments. Rep. Int. Whal. Comm. 21:40-50.
- IWC. 1992. Report of the comprehensive assessment special meeting on North Atlantic fin whales. Rep. Int. Whal. Commn 42:595-644.
- Jennings, S., J.K. Pinnegar, N.V.C. Polunin, K.J. Warr. 2001. Impacts of trawling disturbance on the trophic structure of benthic invertebrate communities. Marine Ecology Progress Series 213: 127-142.
- Johnson, D.L., W.W. Morse, P.L. Berrien, and J.J. Vitaliano. 1999. Essential fish habitat source document: yellowtail flounder, *Limanda ferruginea*, life history and habitat characteristics.
- Kaiser, M.J., J.S. Collie, S.J. Hall, S. Jennings and I.R. Poiner. 2002. Modifications of marine habitats by trawling activities: prognosis and solutions. Fish and Fisheries 3: 114-136.
- Kaiser, M.J., K. Ramsay, C.A. Richardson, F.E. Spence, and A.R. Brand. 2000a. Chronic fishing disturbance has changed shelf sea benthic community structure. Journal of Animal Ecology 69: 494-503.
- Kaiser, M.J., B.E. Spencer, and P.J. Hart. 2000b. Fishing-gear restrictions and conservation of benthic habitat complexity. Conservation Biology 14(5):1512-1525.
- Kaiser, M.J., S.I. Rogers and J.R. Ellis. 1999. Importance of benthic habitat complexity for demersal fish assemblages. In L.R. Benaka, editor. Fish Habitat: Essential fish habitat and rehabilitation. American Fisheries Society, Symposium 22, Bethesda, Maryland.
- Kaiser, M.J., A.S. Hill, K. Ramsay, B.E. Spencer, A.R. Brand, L.O. Veale, K. Prudden, E.I.S. Rees, B.W. Munday, B. Ball, and S.J. Hawkins. 1996a. Benthic disturbance by fishing gear in the Irish Sea: a comparison of beam trawling and scallop dredging. Aquatic Conservation: Marine and Freshwater Ecosystems 6:269-285.
- Kaiser, M.J., D.B. Edwards and B.E. Spencer. 1996b. Infaunal community changes as a result of commercial clam cultivation and harvesting. Aquatic Living Resources 9:57-63.
- Katona, S.K., and J.A. Beard. 1990. Population size, migrations, and feeding aggregations of the humpback whale (*Megaptera novaeangliae*) in the Western North Atlantic Ocean. Rep. Int. Whal. Comm., Special Issue 12: 295-306.
- Kearney/Centaur. 1986. Economic impact of the commercial fishing industry in the United States. Prepared for the National Marine Fisheries Service, Alexandria, Virginia.
- Keinath, J.A., J.A. Musick, and R.A. Byles. 1987. Aspects of the biology of Virginia's sea turtles: 1979-1986. Virginia J. Sci. 38(4):329-336.
- Kelley, J.T. 1998. Mapping the surficial geology of the western Gulf of Maine. Pp. 15-19 in: Northeast Multispecies Amendment 13 SEIS I-1070
December 18, 2003

Effects of fishing gear on the sea floor of New England, E.M. Dorsey and J. Pederson (eds.). MIT Sea Grant Pub. 98-4.

- Kenchington, E.L.R., J. Prena, K.D. Gilkinson, D.C. Gordon Jr., K. MacIssace, C. Bourbonnais, P.J. Schwinghamer, T.W. Rowell, D.L. McKeown, and W.P. Vass. 2001. Effects of experimental otter trawling on the macrofauna of a sandy bottom ecosystem on the Grand Banks of Newfoundland. *Canadian Journal of Fisheries and Aquatic Science* 58:1043-1057.
- Kennedy, P. 1979. *Guide to econometrics*. Cambridge, MA: MIT Press.
- Kenney, R.D. 2000. Are right whales starving? Electronic newsletter of the Center for Coastal Center, posted at www.coastalstudies.org/entanglementupdate/kenney1.html on November 29, 2000. 5pp.
- Kenney, R.D., M.A.M. Hyman, R.E. Owen, G.P. Scott, and H.E. Winn. 1986. Estimation of prey densities required by Western North Atlantic right whales. *Mar. Mamm. Sci.* 2(1): 1-13.
- Kenney, R.D., H.E. Winn, and M.C. Macauley. 1995. Cetaceans in the Great South Channel, 1979-1989: right whale (*Eubalaena glacialis*). *Cont. Shelf Res.* 15:385-414.
- Kitts, A. and E. Thunberg. 1998. Description and Impacts of Northeast Groundfish Fishery Buyout Programs. NOAA/NMFS/NEFSC Reference Document 98-12.
- Klumov, S.K. 1962. The right whale in the Pacific Ocean. In P.I. Usachev (Editor), *Biological marine studies*. *Trud. Inst. Okeanogr.* 58: 202-297.
- Kjesbu, O.S., P. Solemdal, P. Bratland, and M. Fonn. 1996. Variation in annual egg production in individual captive Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences* 53: 610-620.
- Klein-MacPhee, G. 2002a. Righteye Flounders. Family Pleuronictidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Klein-MacPhee, G. 2002b. Cods. Family Gadidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Klein-MacPhee, G. 2002c. Silver Hakes. Family Merlucciidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Klein-MacPhee, G. 2002d. Family Scophthalmidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Klein-MacPhee, G. 2002e. Sea Basses. Family Serranidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Klein-MacPhee, G. 2002f. Porgies. Family Sparidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.

- Klein-MacPhee, G. and B.B. Collette. 2002a. Eelpouts. Family Zoarcidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Klein-MacPhee, G. and B.B. Collette. 2002b. Scorpionfishes. Family Scorpaenida. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Knowlton, A. R., J. Sigurjonsson, J.N. Ciano, and S.D. Kraus. 1992. Long-distance movements of North Atlantic right whales (*Eubalaena glacialis*). *Mar. Mamm. Sci.* 8(4): 397-405.
- Knowlton, A.R., S.D. Kraus, and R.D. Kenney. 1994. Reproduction in North Atlantic right whales (*Eubalaena glacialis*). *Can. J. Zool.* 72: 1297-1305.
- Knutsen, G. M. and S. Tisleth. 1985. Growth, development, and feeding success of Atlantic cod larvae (*Gadus morhua*) related to egg size. *Transactions of the American Fisheries Society* 114: 507-511.
- Kraus, S.D. 1990. Rates and Potential Causes of Mortality in North Atlantic Right Whales (*Eubalaena glacialis*). *Mar. Mamm. Sci.* 6(4):278-291.
- Kreiger, K.J. 2001. Coral (Primnoa) impacted by fishing gear in the Gulf of Alaska. *Proceedings of the First International Symposium on Deep-Sea Corals*. (Also in press in reviewed issue of *Hydrobiologia*).
- Kurkul, Pat. 2000. Essential fish habitat consultation for experimental fishing permit: use of tuna purse seine gear in multispecies Closed Area I on Georges Bank. National Marine Fisheries Service memo to file, September 14, 2000.
- Kurkul, Pat. 2003. Northeast Region, National Marine Fisheries Service. Letter to Thomas Hill, New England Fishery Management Council. July 21, 2003.
- Lallemand, P. J.M. Gates, J. Dirlam, and J. Cho. 1999. The costs of large trawlers in the Northeast. Department of Environmental and Natural Resource Economics, The University of Rhode Island.
- Lallemand, P., J.M. Gates, J. Dirlam, and J. Cho. 1998. The costs of small trawlers in the Northeast. Department of Environmental and Natural Resource Economics, The University of Rhode Island.
- Langan, R.W. 1998. The effect of dredge harvesting on eastern oysters and the associated benthic community. Pp. 108-110 in E.M. Dorsey and J. Pederson, editors. *Effect of Fishing Gear on the Sea Floor of New England*. Conservation Law Foundation. Boston, Massachusetts. 160 p.
- Langton, R.W. and W.E. Robinson. 1990. Faunal associations on scallop grounds in the western Gulf of Maine. *Journal of Experimental Marine Biology and Ecology* 144:157-171.
- Langton, R.W. and W.E. Robinson. 1988. Ecology of the sea scallop, *Placopecten magellanicus* (Gmelin, 1791) in the Gulf of Maine, U.S.A. – a preliminary report. Pp. 243-255 in: *Benthic productivity and marine resources of the Gulf of Maine*, I. Babb and M. Deluca (eds.),

National Undersea Research Program Research Report 88-3, U.S. Department of Commerce, Washington, D.C.

Larson P.F. and R.M. Lee. 1978. Observations on the abundance, distribution and growth of post-larval sea scallops, *Placopecten magellanicus* from the Gulf of Maine. Mar. Ecol. Prog. Ser. 37: 19-25.

Leatherwood, S., and R.R. Reeves. 1983. The Sierra Club handbook of whales and dolphins. Sierra Club Books, San Francisco, California. 302 pp.

Lenihan, H.S. and C.H. Peterson. 1998. How habitat degradation through fishery disturbance enhances impacts of hypoxia on oyster reefs. Ecological Applications 8(1):128-140.

Lindholm, J. and P.J. Auster. 2000. A comparison of structural elements of sand habitats inside and outside of Closed Area II on Georges Bank. Final Report for NOAA Contract 40EQNF900061, Groton, Connecticut. 1-19.

Lindholm, J.B., P.J. Auster, M. Ruth, and L. Kaufman. 2001. Juvenile fish responses to variations in seafloor habitats: modeling the effects of fishing and implications for the design of marine protected areas. Conservation Biology. 15:424-437.

Link, J.S., L.P. Garrison, and F.P. Almeida. 2002a. Ecological interactions between elasmobranchs and groundfish species on the Northeastern U.S. continental shelf. I. Evaluating predation. North American Journal of Fisheries Management 22:550-562.

Link, J.S., J.K.T. Brodziak (eds.). 2002b. Status of the Northeast U. S. Continental Shelf Ecosystem, A report of the Northeast Fisheries Science Center's Ecosystem Status Working Group. Northeast Fisheries Science Center Reference Document 02-11.

Lough, R.G. and D.C. Potter. 1993. Vertical distribution patterns and diel migrations of larval and juvenile haddock *Melanogrammus aeglefinus* and Atlantic cod *Gadus morhua* on Georges Bank. U.S. Fisheries Bulletin 91(2):281-303.

Lutcavage, M. and J.A. Musick. 1985. Aspects of the biology of sea turtles in Virginia. Copeia 1985(2): 449-456.

MacKenzie, C.L., Jr. 1982. Compatibility of invertebrate populations and commercial fishing for ocean quahogs. North American Journal of Fisheries Management 2:270-275.

Maier, P.P., P.H. Wendt, W.A. Roumillat, G.H. Steele, M.V. Levisen, and R. Van Dolah. 1995. Effects of subtidal mechanical clam harvesting on tidal creeks. South Carolina Department of Natural Resources, Marine Resources Division. Final Report. 38 p.

Malik, S., M. W. Brown, S.D. Kraus and B. N. White. 2000. Analysis of mitochondrial DNA diversity within and between North and South Atlantic right whales. Mar. Mammal Sci. 16:545-558.

Manderson, J.P., B.A. Phelan, A.W. Stoner and J. Hilbert. 2000. Predator-prey relations between age-1+ summer flounder and age-0 winter flounder: predator diets, prey selection, and effects of sediments and macrophytes. Journal of Experimental Marine Biology and Ecology 251(1):17-39.

Marine Policy Center, Woods Hole Oceanographic Institution. 2000. Development of an input-

output model for social economic impact assessment of fisheries regulations in New England. MARFIN Final Project Report, National Marine Fisheries Service, Grant Number: NA87FF0548.

- Market Decisions, Inc. 2002. DMR Groundfish Regulation Impact Survey. Market Decisions, Inc.
- Mate, B.M., S.L. Nieuwkerk, and S.D. Kraus. 1997. Satellite monitored movements of the North Atlantic right whale. *J. Wildl. Manage.* 61:1393-1405.
- MacCall, Alec D. 1990. Dynamic Geography of marine fish populations. University of Washington Press, Seattle, WA. 153 pp.
- Mayer, L.M., D.F. Schick, R.H. Findlay and D.L. Rice. 1991. Effects of commercial dragging on sedimentary organic matter. *Marine Environmental Research* 31:249-261.
- Mayo, C.A., and M.K. Marx. 1990. Surface foraging behavior of the North Atlantic right whale, *Eubalaena glacialis*, and associated zooplankton characteristics. *Can. J. Zool.* 68:2214-2220.
- McBride, M. M., L. O'Brien, W. Overholtz, M. Pennington, G. Power, M. Silverman, M. Terceiro, and P. Wood. 1992. Examination of sea sampling data for specified differences in catch rate. Report for the New England Fishery Management Council Groundfish Plan Development Team. Unpublished manuscript.
- McCay, Bonnie and Marie Cieri. 2000. Fishing Ports of the Mid-Atlantic. Report to the Mid-Atlantic Fishery Management Council. Dover, Delaware.
- McCay, Bonnie J., and J. O'Neil. 1998. Social and Economic Characteristics of the Maine Party and Charter Boat Industry. The Ecopolicy Center for Agriculture, the Environment, and Resource Issues, Rutgers University. East Brunswick, New Jersey.
- McCay, Bonnie J., J. O'Neil, J. Velchek. 1999. The New Jersey Party and Charter Boat Industry in 1997: Economics, Job Satisfaction, and Issues. Department of Human Ecology, Rutgers University. East Brunswick, New Jersey.
- McCay, Bonnie J., J. O'Neil, M. Malchoff, and J. Velcheck. 1999. Social and Economic Characteristics of the New York Saltwater Party and Charter Boat Industry, 1997. Department of Human Ecology, Rutgers University. East Brunswick, New Jersey.
- McCay, Bonnie J., Carolyn F. Creed, Alan Christopher Finlayson, Richard Apostle, and Knut Mikalsen. 1995. "Individual Transferable Quotas (ITQs) in Canadian and US Fisheries," *Ocean and Coastal Management* 28(1-3): 85-115.
- McCay, Bonnie J. 1995. "Social and Ecological Implications of ITQs: An Overview," *Ocean and Coastal Management* 28(1-3): 3-22.
- McConnaughey, R.A., K.L. Mier and C.B. Dew. 2000. An examination of chronic trawling on soft bottom benthos of the eastern Bering Sea. *ICES Journal of marine Science* 57:1388-2000.
- McEachran, J.D. 2002. Skates. Family Rajidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- McLoughlin, R.J., P.C. Young, R.B. Martin, and J. Parslow. 1991. The Australian scallop

- dredge: estimates of catching efficiency and associated indirect fishing mortality. Fisheries Research, 11:1-24.
- McMillan, D.G. and W.W. Morse. 1999. Essential fish habitat source document: spiny dogfish *Squalus acanthias*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-150. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 19 p.
- Medcof, J.C. and J.F. Caddy. 1971. Underwater observations on the performance of clam dredges of three types. ICES CM 1971/B:10.
- Meyer, T.L., R.A. Cooper and K.J. Pecci. 1981. The performance and environmental effects of a hydraulic clam dredge. Marine Fisheries Review 43(9):14-22.
- Meylan, A., B. Schroeder, and A. Mosier. 1995. Sea turtle nesting activity in the state of Florida. Fla. Mar. Res. Publ. 52:1-51.
- Milton, S.L., S. Leone-Kabler, A.A. Schulman. And P.L. Lutz. 1994. Effects of hurricane Andrew on the sea turtle nesting beaches of South Florida. Bull. Mar. Sci., 54(3):974-981.
- Minami, Michael. 2000. Using ArcMap. Environmental Systems Research Institute, Inc., Redlands, CA. 528 p.
- Minnesota IMPLAN Group, Inc. 1997. IMPLAN professional: social accounting and impact analysis software. Minnesota IMPLAN Group, Inc., Minneapolis.
- Mirarchi, F. 1998. Bottom trawling on soft substrates. Pp. 80-84 in: Effects of Fishing Gear on the Sea Floor of New England, E.L. Dorsey and J. Pederson (eds.). Conservation Law Foundation, Boston, Massachusetts.
- Mitchell, E. 1974. Present status of the northwest Atlantic fin and other whale stocks. Pages 108-169 in W.E. Schevill (ed) The whale problem: A status report. Harvard University Press, Cambridge, Massachusetts, 419 pp.
- Mitchell, E. and D.G. Chapman. 1977. Preliminary assessment of stocks of northwest Atlantic sei whales (*Balaenoptera borealis*). Rep. Int. Whal. Comm. Special Edition 1:117-120.
- Mizroch, S.A. and A.E. York. 1984. Have pregnancy rates of Southern Hemisphere fin whales, *Balaenoptera physalus*, increased? Rep. Int. Whal. Commn (Spec. Iss. 6):401-410.
- Mooney-Seus, M. and J.E. Dianto. 2000. Beyond Our Shores: Catching Fish in New England Waters Chelsea, MA: Shawmut Printing, 95.
- Moran, M.J. and P.C. Stephenson. 2000. Effects of otter trawling on macrobenthos and management of demersal scalefish fisheries on the continental shelf of north-western Australia. ICES Journal of Marine Science 57:510-516.
- Morgan, Gary R. 1997. Individual quota management in fisheries: Methodologies for determining catch quotas and initial quota allocation. Food and Agriculture Organization of the United Nations. Fisheries Technical Paper 371.

- Morse, W.W., D.L. Johnson, P.L. Berrien, and S.J. Wilk. 1999. Essential fish habitat source document: silver hake *Merluccius bilinearis*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-135. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 42 p.
- Mountain, D.G., R.W. Langton, and L. Watling. 1994. Oceanic processes and benthic substrates: influences on demersal fish habitats and benthic communities. Pp. 20-25 in: Langton, R.W., J.B. Pearce, and J.A. Gibson (eds). Selected Living Resources, Habitat Conditions, and Human Perturbations of the Gulf of Maine: Environmental and Ecological Considerations for Fishery Management. NOAA Technical Memorandum NMFS-NE-106, Woods Hole, MA., 70 p.
- Munroe, T.A. 2002. Herrings. Family Clupeidae. In: B.B. Collette and G. Klein-MacPhee, eds. *Bigelow and Schroeders's Fishes of the Gulf of Maine*. 3rd Edition. Smithsonian Institution Press, Washington, D.C. 748p.
- Murawski, S.A. and F.M. Serchuk. 1989. Environmental effects of offshore dredge fisheries for bivalves. ICES CM 1989/K:27. 12 p.
- Murawski, S. A. 1994. Factors influencing by-catch and discard rates: analyses from multispecies/multifishery sea sampling. *J. Northw. Atl. Fish. Sci.* 19: 31-39.
- Murison, L.D., and D.E. Gaskin. 1989. The distribution of right whales and zooplankton in the Bay of Fundy, Canada. *Can. J. Zool.* 67:1411-1420.
- Murphy, T.M. and S.R. Hopkins. 1984. Aerial and ground surveys of marine turtle nesting beaches in the southeast region. U.S. Final Rept. to NMFS-SEFSC. 73pp.
- Musick, J.A. and C.J. Limpus. 1997. Habitat utilization and migration in juvenile sea turtles. Pp 137-164 In: Lutz, P.L. and J.A. Musick, eds., *The Biology of Sea Turtles*. CRC Press, New York. 432 pp.
- National Marine Fisheries Service (NMFS). 2002. Environmental Assessment for Interim Action. Gloucester, MA.
- National Marine Fisheries Service. 2002. Interim Action to Implement Measures to Reduce Overfishing of the Northeast Fishery Complex Under the Northeast Multispecies Fishery Management Plan.
- National Marine Fisheries Service. 2001a. Report to Congress: Report on the Status of Fisheries of the United States.
- National Marine Fisheries Service. 2001b. Draft Environmental Assessment Incorporating a Regulatory Impact Review for an Experimental Fishery to Allow Tuna Purse Seine Vessels to Fish in the Northeast Multispecies Closed Area I on Georges Bank.
- National Marine Fisheries Service. 1999. Atlantic Highly Migratory Species Fishery Management Plan.
- NMFS. 1998a. Recovery plan for the shortnose sturgeon (*Acipenser brevirostrum*). Prepared by the Shortnose Sturgeon Recovery Team for the National Marine Fisheries Service, Silver Spring, Maryland. 104 pages.
- NMFS. 1998b. Draft recovery plans for the fin whale (*Balaenoptera physalus*) and sei whale (*Balaenoptera borealis*). Prepared by R.R. Reeves, G.K. Silber, and P.M. Payne for the National Marine Fisheries

- Service, Silver Spring, Maryland. July 1998.
- NMFS. 1998c. Recovery plan for the blue whale (*Balaenoptera musculus*). Prepared by Reeves, R.R., P.J. Clapham, and R.L. Brownell, Jr. for the National Marine Fisheries Service, Silver Spring, Maryland. Mitchell, E. 1974. Present status of the northwest Atlantic fin and other whale stocks. Pages 108-169 in W. E. Schevill (ed) *The Whale Problem: A status report*. Harvard University Press. Cambridge, Massachusetts, 419pp.
- NMFS. 1991a. Final recovery plan for the humpback whale (*Megaptera novaeangliae*). Prepared by the Humpback Whale Recovery Team for the National Marine Fisheries Service, Silver Spring, Maryland. 105 pp.
- NMFS. 1991b. Final recovery plan for the North Atlantic right whale (*Eubalaena glacialis*). Prepared by the Right Whale Recovery Team for the National Marine Fisheries Service, Silver Spring, Maryland. 86 pp.
- National Marine Fisheries Service and U.S. Fish and Wildlife Service. 1995. Status reviews for sea turtles listed under the Endangered Species Act of 1973. Silver Spring, MD: National Marine Fisheries Service, 139.
- NMFS and USFWS. 1995. Status reviews for sea turtles listed under the Endangered Species Act of 1973. National Marine Fisheries Service, Silver Spring, Maryland. 139 pp.
- NMFS and USFWS. 1992. Recovery plan for leatherback turtles in the U.S. Caribbean, Atlantic, and Gulf of Mexico. National Marine Fisheries Service, Washington, D.C. 65 pp.
- NMFS and USFWS. 1991a. Recovery plan for U.S. population of Atlantic green turtle. National Marine Fisheries Service, Washington, D.C. 52 pp.
- NMFS and USFWS. 1991b. Recovery plan for the U.S. population of loggerhead turtle. National Marine Fisheries Service, Washington, D.C. 64 p.
- National Oceanic and Atmospheric Administration-National Marine Fisheries Service. March 2001. Guidance for Social Impact Assessment. Draft memorandum distributed to technical staff of fishery management councils, 38.
- National Research Council. 2002. Effects of Trawling and Dredging on Seafloor Habitat. National Academy Press. 126 p.
- New England Fishery Management Council (NEFMC). 2002. FMP for deep-sea red crab *Chaceon quinque-dens*. Volume I.
- New England Fishery Management Council Staff. 2001a. Report from Groundfish Social Impact Informational Meetings.
- New England Fishery Management Council Staff. 2001b. Summary Statistics for Gross Revenues and Landings of Multispecies Permit Holders since FY 1994.
- New England Fishery Management Council. 2001c. Report of the Multispecies Monitoring Committee.
- New England Fishery Management Council. 2000a. Amendment 12 to the Northeast Multispecies Fishery Management Plan.

- New England Fishery Management Council. 2000b. Framework Adjustment 33 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council. 2000c. Report of the Multispecies Monitoring Committee.
- New England Fishery Management Council. 1999a. Amendment 9 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council. 1999b. Framework Adjustment 26 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council. 1999c. Framework Adjustment 27 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council. 1999d. Framework Adjustment 30 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council. 1998a. Framework Adjustment 25 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council. 1998b. Amendment 9 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council. 1998c. Omnibus Essential Fish Habitat Amendment (Amendment 11 to the Northeast Multispecies FMP, Amendment 9 to the Sea Scallop FMP, Amendment 1 to the Monkfish FMP, Amendment 1 to the Atlantic Salmon FMP, and Sections of the Atlantic Herring FMP).
- New England Fishery Management Council. 1996. Amendment 7 to the Northeast Multispecies Fishery Management Plan.
- New England Fishery Management Council. 1994. Amendment 5 to the Northeast Multispecies Fishery Management Plan.
- NOAA Tech. Mem. NMFS-NE-140. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 29 p.
- Northeast Fisheries Science Center (NEFSC). 2002a. 19 March 2002. Final Report of the Working Group on Re-Evaluation of Biological Reference Points for New England Groundfish. NMFS-NEFSC. Woods Hole, MA
- Northeast Fisheries Science Center (NEFSC). 2002b. Assessment of 20 Northeast Groundfish Stocks through 2001: a report of the Groundfish Assessment Review Meeting (GARM), Northeast fisheries Science Center, Woods Hole, MA, October 8-11, 2001.
- Northeast Fisheries Science Center (NEFSC). 2002c. Report of the 35th Northeast Regional Stock Assessment Workshop (35th SAW): Consensus Summary of Assessments. NEFSC Reference Document 02-014).
- Northeast Fisheries Science Center. 2001a. 32nd Northeast Regional Stock Assessment

- Workshop (32nd SAW): Stock Assessment Review Committee (SARC) Consensus Summary of Assessments. (NEFSC Ref. Doc. 01-05).
- Northeast Fisheries Science Center. 2001b. 33rd Northeast Regional Stock Assessment Workshop (33rd SAW): Stock Assessment Review Committee (SARC) Advisory Report on Stock Status. NEFSC Ref. Doc. 01-19.
- Northeast Fisheries Science Center. 2001c. TRAC Advisory Report on Stock Status: A report of the Fourth Meeting of the Transboundary Resources Assessment Committee (TRAC). St. Andrews Biological Station, St. Andrews, New Brunswick. April 17-20, 2001. (NEFSC Ref. Doc. 01-08).
- Northeast Fisheries Science Center. 2001d. 33rd Stock Northeast Regional Stock Assessment Workshop (33rd SAW): Stock Assessment Review Committee (SARC) Consensus Summary of Assessments. (NEFSC Ref. Doc. 01-18).
- Northeast Fisheries Science Center. 2000. Assessment of 19 Northeast Groundfish Stocks through 2000. Report of the SAW Northern Demersal and Southern Demersal Working Groups. Submitted to the NEFMC Multi-Species Monitoring Committee. Inter-Sessional meeting, August, 2000.
- Northeast Fisheries Science Center. 2000. 30th Stock assessment workshop report. Woods hole, MA. April 2000. NMFS-NEFSC Ref. Doc. 00-03.
- Northeast Region Essential Fish Habitat Steering Committee (NREFHSC). 2002. Workshop on the Effects of Fishing Gear on Marine Habitats Off the Northeastern United States, October 23-25, 2001, Boston, Massachusetts. Northeast Fish Sci Cent Ref Doc 02-01; 86.
- O'Brien, Loretta. 1999. Factors influencing the rate of sexual maturity and the effect on spawning stock for Georges Bank and Gulf of Maine Atlantic cod (*Gadus morhua*) stocks. Journal of Northwest Atlantic Fisheries Science 25: 179-203.
- O'Brien, Loretta, Jay Burnett, Ralph K. Mayo. 1993. Maturation of nineteen species of finfish off the Northeast Coast of the United States, 1985-1990. U. S. Department of Commerce, NOAA Technical Report NMFS 113.
- Ogren, L.H. 1988. Biology and ecology of sea turtles. Prepared for the National Marine Fisheries Service, Panama City Laboratory. Sept 7.
- Ohman, M.C., A. Rajasuriya, and O. Linden. 1993. Human disturbance on coral reefs in Sri Lanka: a case study. *Ambio* 22: 474-480.
- Orth, R.J., K.A. Moore, D.J. Wilcox and J.R. Fishman. 1998. Chincoteague Bay, Virginia: effectiveness of the SAV sanctuary and revegetation of SAV habitat disturbed by clam dredging. Unpublished Report to the Virginia Marine Resources Commission.
- Overholtz, W.J. and A.V. Tyler. 1985. Long-term responses of the demersal fish assemblages of Georges Bank. *U.S. Fisheries Bulletin* 83(4):507-520.
- Packer, D.B., C.A. Zetlin and J.J. Vitaliano. In press (a). Essential fish habitat source document: barndoor skate, *Dipturus laevis*, life history and habitat characteristics. NOAA Tech. Mem. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA.
- Packer, D.B., C.A. Zetlin and J.J. Vitaliano. In press (b). Essential fish habitat source

- document: clearnose skate, *Raja eglanteria*, life history and habitat characteristics. NOAA Tech. Mem. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA.
- Packer, D.B., C.A. Zetlin and J.J. Vitaliano. In press (c). Essential fish habitat source document: little skate, *Leucoraja erinacea*, life history and habitat characteristics. NOAA Tech. Mem. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA.
- Packer, D.B., C.A. Zetlin and J.J. Vitaliano. In press (d). Essential fish habitat source document: rosette skate, *Leucoraja garmani virginica*, life history and habitat characteristics. NOAA Tech. Mem. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA.
- Packer, D.B., C.A. Zetlin and J.J. Vitaliano. In press (e). Essential fish habitat source document: smooth skate, *Malacoraga senta*, life history and habitat characteristics. NOAA Tech. Mem.
- Packer, D.B., C.A. Zetlin and J.J. Vitaliano. In press (f). Essential fish habitat source document: thorny skate, *Amblyraja radiata*, life history and habitat characteristics. NOAA Tech. Mem. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA.
- Packer, D.B., C.A. Zetlin and J.J. Vitaliano. In press (g). Essential fish habitat source document: Winter skate, *Leucoraja ocellata*, life history and habitat characteristics. NOAA Tech. Mem. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA.
- Packer, D.B., L.M. Cargnelli, S.J. Griesbach, and S.E. Shumway. 1999(a). Essential fish habitat source document: Atlantic sea scallop, *Placopecten magellanicus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-134. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 21 p.
- Packer, D.B., S.J. Griesbach, P.L. Berrien, C.A. Zetlin, D.L. Johnson, and W.W. Morse. 1999(b). Essential fish habitat source document: summer flounder, *Paralichthys dentatus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-151. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA.
- Pálsson, Gísli. 1998. "The Virtual Aquarium: Commodity Fiction and Cod Fishing," *Ecological Economics* 24: 275–288.
- Pálsson, Gísli and Agnar Helgason. 1995. "Figuring Fish and Measuring Men: The Quota System in the Icelandic Cod Fishery." *Ocean and Coastal Management* 28(1-3): 117–146.
- Payne, A. I., R. Cook, E. Bell, R. K. Mohn, and M. McAllister. February 3 - 8, 2003. Report on the Groundfish Science Peer Review Meeting. Center for Independent Experts.
- Payne, P.M., D.N. Wiley, S.B. Young, S. Pittman, P.J. Clapham, and J.W. Jossi. 1990. Recent fluctuations in the abundance of baleen whales in the southern Gulf of Maine in relation to changes in selected prey. *Fish. Bull.* 88 (4): 687-696
- Pereira, J.J., R. Goldberg, J.J. Ziskowski, P.L. Berrien, W.W. Morse, and D.L. Johnson. 1999. Essential fish habitat source document: winter flounder, *Pseudopleuronectes americanus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-138. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 39 p.
- Perry, S.L., D.P. DeMaster, and G.K. Silber. 1999. The Sperm Whale In: The great whales: History and status of six species listed as endangered under the U.S. Endangered Species Act of 1973. Mar.

Fish. Rev. Special Edition. 61(1): 59-74.

Petrakis, G., D. N. MacLennan, and A. W. Newton. 2001. Day-night depth effects on catch rates during trawl surveys in the North Sea. *ICES Journal of Marine Science* 58: 50-60.

Phelan, B.A., J.P. Manderson, A.W. Stoner and A.J. Bejda. 2001. Size-related shifts in the habitat associations of YOY winter flounder: field observations and laboratory experiments with sediments and prey. *Journal of Experimental Marine Biology and Ecology* 257(2):297-315.

Pikanowski, R.A., W.W. Morse, P.L. Berrien, D.L. Johnson, D.G. McMillan. 1999. Essential fish habitat source document: redfish, *Sebastes* spp., life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-132. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 19 p.

Pilskaln, C.H., J.H. Churchill, and L.M. Mayer. 1998. Resuspension of sediment by bottom trawling in the Gulf of Maine and potential geochemical consequences. *Conservation Biology* 12(6):1223-1229.

Pinhorn, A. T. and R. G. Halliday. 2001. The regulation of exploitation pattern in North Atlantic groundfish stocks. *Fisheries Research* 53: 25-37.

Pol, Michael and W. Glenn Hovermale. 2000. A discussion of selectivity. Unpublished manuscript, Massachusetts Division of Marine Fisheries.

Pol, Michael. 2003. Massachusetts Division of Marine Fisheries, Conservation Engineering Branch. Personal communication, July 29, 2003.

Pollnac, Richard B. and S.J. Littlefield. 1983. Sociocultural aspects of fisheries management. *Ocean Development and International Law Journal* 12 (3-4): 209-246.

Poppe, L.J., J.S. Schlee, B. Butman, and C.M. Lane. 1989. Map showing distribution of surficial sediment, Gulf of Maine and Georges Bank. U.S. Geological Survey Miscellaneous Investigations Series, Map 1-1986-A.

Portland Fish Exchange. [Online: web] URL: <http://www.portlandfishexchange.com>

Pranovi, F. and O. Giovanardi. 1994. The impact of hydraulic dredging for short-necked clams, *Tapes* spp., on an infaunal community on the lagoon of Venice. *Scientia Marina* 58(4):345-353.

Pratt, S. 1973. Benthic fauna. Pp. 5-1 to 5-70 in: Coastal and offshore environmental inventory, Cape Hatteras to Nantucket Shoals. University of Rhode Island, Marine Publication Series No. 2. Kingston, RI.

Prena, J., P. Schwinghamer, T.W. Rowell, D.C. Gordon, Jr., K.D. Gilkinson, W.P. Vass, and D.L. McKeown. 1999. Experimental otter trawling on a sandy bottom ecosystem of the Grand Banks of Newfoundland: analysis of trawl bycatch and effect on epifauna. *Marine Ecology Progress Series* 181:107-124.

Prescott, R.L. 1988. Leatherbacks in Cape Cod Bay, Massachusetts, 1977-1987, p 83-84 In: B.A. Schroeder (comp.), Proceedings of the Eighth Annual Workshop on Sea Turtle Conservation and Biology. NOAA Technical Memorandum NMFS-SEFC-214.

Pritchard, P.C.H. 1969. Endangered species: Kemp's ridley turtle. *Florida Naturalist*, 49:15-19.

- Pritchard, P.C.H. 1982. Nesting of the leatherback turtle, *Dermochelys coriacea*, in Pacific, Mexico, with a new estimate of the world population status. *Copeia* 1982:741-747.
- Rebel, T.P. 1974. Sea turtles and the turtle industry of the West Indies, Florida and the Gulf of Mexico. Univ. Miami Press, Coral Gables, Florida.
- Reeves, R.R., and Mitchell, E. 1988. History of whaling in and near North Carolina. NOAA Tech. Rep. NMFS 65: 28 pp.
- Reeves, R.R., Breiwick, J.M., and Mitchell, E. 1992. Pre-exploitation abundance of right whales off the eastern United States. Pp. 5-7 in J. Hain (ed.), The right whale in the western North Atlantic: a science and management workshop, 14-15 April 1992, Silver Spring, Maryland. National Marine Fisheries Service, NEFSC Ref. Doc. 92-05.
- Reid, R.N., L.M. Cargnelli, S.J. Griesbach, D.B. Packer, D.L. Johnson, C.A. Zetline, W.W. Morse and P.L. Berrien. 1999. Essential fish habitat source document: Atlantic herring, *Clupea harengus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-126. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 48 p.
- Reid, R.N. and F.W. Steimle. 1988. Pp. 125-160 in: A.L. Pacheco (ed.), Characterization of the middle Atlantic water management unit of the northeast regional action plan. NOAA Technical Memorandum NMFS-F/NEC-56. Woods Hole, MA., 322 p.
- Reijnders, P.J.H. and A. Aguilar. 2002. Pollution and Marine Mammals. *In*: W.F. Perrin, B. Würsig, J.G.M. Thewissen (eds.) Encyclopedia of Marine Mammals. Academic Press. p. 948-957.
- Reimann, B. and E. Hoffman. 1991. Ecological consequences of dredging and bottom trawling in the Limfjord, Denmark. *Marine Ecology Progress Series* 69:171-178.
- Reise, K. 1982. Long term changes in the macrobenthic invertebrate fauna of the Wadden Sea: Are polychaetes about to take over? *Netherlands Journal of Sea Research* 16:29-36.
- Reise, K. and A. Schubert. 1987. Macrobenthic turnover in the subtidal Wadden Sea: the Norderaue revisited after 60 years. *Helgolander Meeresunters* 41:69-82.
- Riesen, W. and K. Reise 1982. Macrobenthos of the subtidal Wadden Sea: revisited after 55 years. *Helgolander Meeresunters* 35:409-423.
- Robbins, J., and D. Mattila. 1999. Monitoring entanglement scars on the caudal peduncle of Gulf of Maine humpback whales. Report to the National Marine Fisheries Service. Order No. 40EANF800288. 15 pp.
- Roheim, C. in prep. Market information and fisheries management: improving efficiency in Northeastern U.S. groundfish management systems.
- Root, R.B. 1967. The niche exploitation pattern of the blue-gray gnatcatcher. *Ecol Monogr* 37:317-350.
- Rosenbaum, H.C., M.G. Egan, P.J. Clapham, R.L. Brownell Jr., S.Malik, M. Brown, B. White, P. Walsh and R.DeSalle. 2000. Assessing a century of genetic change in North Atlantic right whales

(*Eubalaena glacialis*). Cons. Biol.

- Ross, J.P. 1979. Green turtle, *Chelonia mydas*, Background paper, summary of the status of sea turtles. Report to WWF/IUCN. 4pp.
- Rothschild, Brian R. and Wendell Brown. 2002. A High-Resolution, Industry-Conducted, Fishery Resource Survey: A Cooperative Project Involving The School for Marine Science and Technology/University of Massachusetts Dartmouth; Massachusetts Fisheries Recovery Commission; Trawler Survival Fund; and Massachusetts Division of Marine Fisheries. SMAST/University of Massachusetts - Dartmouth.
- Rountree, B.P., P. Clay, S. Steinbeck, and J. Walden. 2001. Status of the Fishery Resources off the Northeastern United States. NOAA/NMFS/NEFSC, Woods Hole, MA.
- Rountree, R.A. and K.W. Able. 1992. Foraging habits, growth, and temporal patterns of salt-marsh creek habitat use by young-of-the-year summer flounder in New Jersey. Transactions American Fisheries Society 121(6):765-776.
- Ruais, Rich. 2000. East Coast Tuna Association. Personal communication.
- Sainsbury, J.C. 1996. *Commercial Fishing methods: An introduction to vessels and gears*, 3rd Ed. Fishing News Books, Oxford, England.
- Sainsbury, K.J., R.A. Campbell, R. Lindholm and A.W. Whitlaw. 1997. Experimental management of an Australian multispecies fishery: examining the possibility of trawl-induced habitat modification. In Pikitch, E.K., D.D. Huppert and M.P. Sissenwine, eds. Global Trends: Fisheries management. American Fisheries Society, Symposium 20, Bethesda, Maryland.
- Salz, R. J., D. K. Loomis, M. R. Ross, and S. Steinback. Unpublished. A Baseline Socio-Economic Study of Massachusetts' Marine Recreational Fisheries. NOAA Technical Memorandum NMFS-NE-(forthcoming). National Marine Fisheries Service, Northeast Fisheries Science Center. Woods Hole, MA.
- Sanchez, P., M. Demestre, M. Ramon, and M.J. Kaiser. 2000. The impact of otter trawling on mud communities in the northwestern Mediterranean. ICES Journal of Marine Science 57:1352-1358.
- Schaeff, C.M., Kraus, S.D., Brown, M.W., Perkins, J.S., Payne, R., and White, B.N. 1997. Comparison of genetic variability of North and South Atlantic right whales (*Eubalaena*), using DNA fingerprinting. Can. J. Zool. 75:1073-1080.
- Schevill, W.E., W.A. Watkins, and K.E. Moore. 1986. Status of *Eubalaena glacialis* off Cape Cod. Rep. Int. Whal. Comm., Special Issue 10: 79-82.
- Schick, Dan. Maine Department of Marine Resources.
- Schmitz, W.J., W.R. Wright, and N.G. Hogg. 1987. Physical oceanography. Pp. 27-56 in: J.D. Milliman and W.R. Wright (eds.), The marine environment of the U.S. Atlantic continental slope and rise. Jones and Bartlett Publishers, Inc., Boston, MA.
- Schwinghamer, P., D.C. Gordon, Jr., T.W. Rowell, J. Prena, D.L. McKeown, G. Sonnichsen, and

- J.Y. Guigne. 1998. Effects of experimental otter trawling on surficial sediment properties of a sandy-bottom ecosystem of the Grand Banks of Newfoundland. *Conservation Biology* 12(6):1215-1222.
- Scida, Pasquale. June 29, 2001. Northeast Regional Office, National Marine Fisheries Service. Gloucester, MA. Personal communication.
- Seipt, I., P.J. Clapham, C.A. Mayo, and M.P. Hawvermale. 1990. Population characteristics of individually identified fin whales, *Balaenoptera physalus*, in Massachusetts Bay. *Fish. Bull.* 88:271-278.
- Shepard, F.P., N.F. Marshall, P.A. McLonghlin, and F.G. Sullivan. 1979. Currents in submarine canyons and other sea valleys. *American Association of Petroleum and Geology, Studies in Geology* No. 8.
- Shepherd, G. and J. Forrester. 1987. Diurnal variation in catchability during bottom trawl surveys of the northeastern United States. *International Council for the Exploration of the Seas C.M./B:44.*
- Sherman, K.J., N.A. Jaworski, T.J. Smayda (eds). 1996. *The Northeast Shelf Ecosystem – Assessment, Sustainability, and Management.* Blackwell Science, Inc. Cambridge, MA.
- Shoop, C.R. and R.D. Kenney. 1992. Seasonal distributions and abundance of loggerhead and leatherback sea turtles in waters of the northeastern United States. *Herpetol. Monogr.* 6: 43-67. Rebel, T.P. 1974. *Sea turtles and the turtle industry of the West Indies, Florida and the Gulf of Mexico.* Univ. Miami Press, Coral Gables, Florida.
- Skrobe, L. G., Beutel, D. L., Castor, K., Valliere, A., Gibson, M., Lazar, N., Borden, D., Lee, L. 2003. Southern New England Yellowtail Flounder Size Selectivity for 6.0" diamond, 6.5" square, 6.5" diamond, and 7.0" square shaped mesh codends. Technical Report 1-03, Rhode Island Gear Conservation Engineering Working Group, University of Rhode Island Sea Grant.
- Smith, E., M.A. Alexander, M.M. Blake, L. Gunn, P.T. Howell, M.W. Johnson, R.E. MacLeod, R.F. Sampson, D.G. Simpson, W.H. Webb, L.L. Stewart, P.J. Auster, N.K. Bender, K. Buchholz, J. Crawford, and T.J. Visel. 1985. A study of lobster fisheries in Connecticut waters of Long Island Sound with special reference to the effects of trawling on lobsters. Unpublished report, Connecticut Department of Environmental Protection, Marine Fisheries Program, Hartford, CT.
- Smolowitz, R. J., P. J. Struhsaker, C. Goudey and H. Kite-Powell. 1997. Result of Gear Modification Test to Reduce Bycatches of Commercial Finfish in Sea Scallop Dredges. Final Report: NMFS/NOAA Grant #NA66FD0026.
- Smolowitz, R. 1998. Bottom tending gear used in New England. In: *Effects of Fishing Gear on the Sea Floor of New England* (E.L. Dorsey and J. Pederson, eds). Conservation Law Foundation, Boston, Massachusetts.
- Southeast Fisheries Science Center. 2001. Stock assessments of loggerheads and leatherback sea turtles and an assessment of the impact of the pelagic longline fishery on the loggerhead and leatherback sea turtles of the Western North Atlantic. U.S. Department of Commerce, National Marine Fisheries Service, Miami, FL, SEFSC Contribution PRD-00/01-08; Parts I-III and Appendices I-IV. NOAA Tech. Memo NMFS-SEFSC-455, 343 pp.
- Sparks-McConkey, P.J. and L. Watling. 2001. Effects on the ecological integrity of a soft-

bottom habitat from a trawling disturbance. *Hydrobiologia* 456:73-85.

Spotila, J.R., A.E. Dunham, A.J. Leslie, A.C. Steyermark, P.T. Plotkin, and F.V. Paladino. 1996. Worldwide Population Decline of *Demochelys coriacea*: Are Leatherback Turtles Going Extinct? *Chelonian Conservation and Biology* 2(2): 209-222.

Steimle, F.W. and C. Zetlin. 2000. Reef habitats in the middle Atlantic bight: abundance, distribution, associated biological communities, and fishery resource use. *Marine Fisheries Review* 62(2):24-42.

Steimle, F.W., C.A. Zetlin and S. Chang. 2001. Essential fish habitat source document: red crab, *Chaceon (Geryon) quinquedens*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-163. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 36 p.

Steimle, F.W., C.A. Zetlin, P.L. Berrien, and S. Chang. 1999a. Essential fish habitat source document: black sea bass, *Centropristis striata*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-143. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 42 p.

Steimle, F.W., C.A. Zetlin, P.L. Berrien, D.L. Johnson and S. Chang. 1999b. Essential fish habitat source document: tilefish, *Lopholatilus chamaeleonticeps*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-152. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 30 p.

Steimle, F.W., W.W. Morse, and D.L. Johnson. 1999c. Essential fish habitat source document: goosefish, *Lophius americanus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-127. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 31 p.

Steimle, F.W., W.W. Morse, P.L. Berrien, D.L. Johnson. 1999d. Essential fish habitat source document: red hake, *Urophycis chuss*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-133. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 34 p.

Steimle, F.W., W.W. Morse, P.L. Berrien, D.L. Johnson, and C.A. Zetlin. 1999e. Essential fish habitat source document: ocean pout, *Macrozoarces americanus*, life history and habitat characteristics. NOAA Tech. Mem. NMFS-NE-129. National Marine Fisheries Service Northeast Fisheries Science Center, Woods Hole, MA. 26 p.

Steinback, S. and B. Gentner. 2001. Marine Angler Expenditures in the Northeast Region, 1998. NOAA Technical Memorandum NMFS-F-SPO-47. NMFS Office of Science and Technology, Fisheries Statistics and Economics Division. Silver Spring, MD.

Steinback, S., and B. Gentner. 2001. Marine angler expenditures in the Northeast region, 1998. NOAA Technical Memorandum NMFS-F/SPO-47.

Steinback, S., J. O'Neil, E. Thunberg, A. Gautam, and M. Osborn. 1999. Volume I: Summary Report of Methods and Descriptive Statistics for the 1994 Northeast Region Marine Recreational Economics Survey. NOAA Technical Memorandum (NMFS-F/SPO-37). NMFS Office of Science and Technology, Fisheries Statistics and Economics Division. Silver Spring, MD.

Stephan, C.D., R.L. Peuser and M.S. Fonseca. 2000. Evaluating fishing gear impacts to Northeast Multispecies Amendment 13 SEIS
December 18, 2003

- submerged aquatic vegetation and determining mitigation strategies. ASMFC Habitat Management Series No. 5. Atlantic States Marine Fisheries Commission, Washington, DC.
- Steves, B.P., R.K. Cowen, and M.H. Malchoff. 1998. Settlement and nursery habitats for demersal fishes on the continental shelf of the New York Bight. U.S. Fisheries Bulletin 98:167-188.
- Stone, S.L., T.A. Lowery, J.D. Field, S.H. Jury, D.M. Nelson, M.E. Monaco, C.D. Williams, and L. Andreasen. 1994. Distribution and abundance of fishes and invertebrates in mid-Atlantic estuaries. ELMR Rep. No. 12. NOAA/NOS Strategic Environmental Assessments Division, Silver Spring, MD 280 p.
- Stoner, A.W., J.P. Manderson and J.P. Pessutti 2001. Spatially explicit analysis of estuarine habitat for juvenile winter flounder: combining GAM and GIS. *Journal of Experimental Marine Biology and Ecology* 213:253-271.
- Stumpf, R.P. and R.B. Biggs. 1988. Surficial morphology and sediments of the continental shelf of the middle Atlantic bight. Pp. 51-72 in: A.L. Pacheco (ed.), *Characterization of the middle Atlantic water management unit of the northeast regional action plan*. NOAA Technical Memorandum NMFS-F/NEC-56. Woods Hole, MA., 322 p.
- Swingle, W.M., S.G. Barco, T.D. Pitchford, W.A. McLellan and D.A. Pabst. 1993. Appearance of juvenile humpback whales feeding in the nearshore waters of Virginia. *Mar. Mammal Sci.* 9:309-315.
- Techlaw Inc. May 2001. *The Economic Contribution of the Sport Fishing, Commercial Fishing, and Seafood Industries to New York State, Final Report*. Funded by and prepared for New York Sea Grant.
- Terwilliger, K. and J.A. Musick. 1995. Virginia sea turtle and marine mammal conservation team. Management plan for sea turtles and marine mammals in Virginia. Final Rept to NOAA, 56 pp.
- Thayer, Pete. Maine Department of Marine Resources.
- Theroux, R.B. and M.D. Grosslein. 1987. Benthic fauna. Pp. 283-195 in: R.H. Backus (ed.), *Georges Bank*. MIT Press, Cambridge, MA.
- Theroux, R.B. and R.L. Wigley. 1998. Quantitative composition and distribution of the macrobenthic invertebrate fauna of the continental shelf ecosystems of the northeastern United States. NOAA Technical Report NMFS 140. U.S. Dept. of Commerce, Seattle, WA.
- Thrush, S.F., J.E. Hewitt, V.J. Cummings, and P.K. Dayton. 1995. The impact of habitat disturbance by scallop dredging on marine benthic communities: What can be predicted from the results of experiments? *Marine Ecology Progress Series* 129(1-3):141-150.
- Thrush, S.F., J.E. Hewitt, V.J. Cummings, P.K. Dayton, M. Cryer, S.J. Turner, G.A. Funnell, R.G. Budd, C.J. Milcurn, and M.R. Wilkinson. 1998. Disturbance of the marine benthic habitat by commercial fishing: impacts at the scale of the fishery. *Ecological Applications* 8(3):866-879.
- Thunberg, E., S. Steinback, G. Gray, A. Gautam, and M. Osborn. 1999. Volume III: Summary Report of Methods and Descriptive Statistics for the 1994 Northeast Region Marine Recreational Fishing Participation Survey. NOAA Technical Memorandum (NMFS-F/SPO-39). NMFS Office of Science and Technology, Fisheries Statistics and Economics Division. Silver Spring, MD.
- Townsend, D.W. 1992. An overview of the oceanography and biological productivity of the Northeast Multispecies Amendment 13 SEIS

- Gulf of Maine. Pp. 5-26 in: Townsend, D.W. and P.F. Larsen (eds). The Gulf of Maine, NOAA Coastal Ocean Program Regional Synthesis Series Number 1. Silver Spring, MD.
- Trippel, E. A. and M. J. Morgan. 1994. Age-specific paternal influences on reproductive success of Atlantic cod (*Gadus morhua* L.) of the Grand Banks, Newfoundland. ICES Mar. Sci. Symp. 198: 414-422.
- Tschernij, Vesa and Rene Holst. 1999. Evidence of factors at vessel-level affecting codend selectivity in Baltic cod demersal trawl fishery. ICES CM 1999/R:02.
- Tucholke, B.E. 1987. Submarine geology. Pp. 56-113 in: J.D. Milliman and W.R. Wright (eds.), The marine environment of the U.S. Atlantic continental slope and rise. Jones and Bartlett Publishers, Inc., Boston, MA.
- Tuck, I.D., N. Bailey, M. Harding, G. Sangster, T. Howell, N. Graham and M. Breen. 2000. The impact of water jet dredging for razor clams, *Ensis* spp., in a shallow sandy subtidal environment. Journal of Sea Research 43:65-81.
- Tuck, I.D., S.J. Hall, M.R. Robertson, E. Armstrong, and D.J. Basford. 1998. Effects of physical trawling disturbance in a previously unfished sheltered Scottish sea loch. Marine Ecology Progress Series 162:227-242.
- Tupper, M. and R. G. Boutilier. 1995. Effects of habitat on settlement, growth, and postsettlement survival of Atlantic cod (*Gadus morhua*). Canadian Journal of Fisheries and Aquatic Science 52:1834-1841.
- Turtle Expert Working Group (TEWG). 1998. An assessment of the Kemp's ridley (*Lepidochelys kempii*) and loggerhead (*Caretta caretta*) sea turtle populations in the Western North Atlantic. NOAA Technical Memorandum NMFS-SEFSC-409. 96 pp.
- Turtle Expert Working Group (TEWG). 2000. Assessment update for the Kemp's ridley and loggerhead sea turtle populations in the western North Atlantic. U.S. Dep. Commer. NOAA Tech. Mem. NMFS-SEFSC-444, 115 pp.
- USFWS. 1997. Synopsis of the biological data on the green turtle, *Chelonia mydas* (Linnaeus 1758). Biological Report 97(1). U.S. Fish and Wildlife Service, Washington, D.C. 120 pp.
- USFWS and NMFS. 1992. Recovery plan for the Kemp's Ridley sea turtle (*Lepidochelys kempii*). National Marine Fisheries Service, St. Petersburg, FL. 40 p.
- Valentine, P. 1998. Brief notes on habitat geology and clay pipe habitat on Stellwagen Bank. In Dorsey, E.M. and J. Pederson, eds. Effects of Fishing Gear on the Sea Floor of New England. Boston, MA: Conservation Law Foundation. 119-120.
- Valentine, P.C. and R.G. Lough. 1991. The sea floor environment and the fishery of eastern Georges bank. Dept. of Interior, U.S. Geological Survey, Open File Report 91-439.
- Valentine, P.C., E.W. Strom, R.G. Lough, and C.L. Brown. 1993. Maps showing the sedimentary environment of eastern Georges bank. U.S. Geological Survey, Miscellaneous Investigations Series, Map I-2279-B, scale 1:250,000.
- Vallin, L. and A. Nissling. 2000. Maternal effects on egg size and egg buoyancy of Baltic cod,

- Gadus morhua*: Implications for stock structure effects on recruitment. Fisheries Research 49: 21-37.
- Van Dolah, R.F., P.H. Wendt, and N. Nicholson. 1987. Effects of a research trawl on a hard-bottom assemblage of sponges and corals. Fisheries Research 5:39-54.
- Veale, L.O., A.S. Hill, S.J. Hawkins, and A.R. Brand. 2000. Effects of long-term physical disturbance by commercial scallop fishing on subtidal epifaunal assemblages and habitats. Marine Biology 137(2):325-337.
- Wallace, D.E. 1997. The molluscan fisheries of Maine. NOAA Tech. Rept. NMFS 127:63-85
- Walsh, H.J., D.S. Peters, and D.P. Cyrus. 1999. Habitat utilization by small flatfishes in a North Carolina estuary. Estuaries 22(3B):803-813.
- Waring, G.T., C.P. Fairfield, C.M. Ruhsam, and M. Sano. 1993. Sperm whales associated with Gulf Stream features off the northeastern USA shelf. Fish. Oceanogr. 2(2):101-105.
- Waring, G. T., J.M. Quintal, S. L. Swartz, eds. 2000. U.S. and Gulf of Mexico marine mammal stock assessments. NOAA Tech. Mem. NMFS-NE-162.
- Waring, G.T., J.M. Quintal, S.L. Swartz (eds). 2001. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments - 2001. NOAA Technical Memorandum NMFS-NE-168.
- Watkins, W.A., and W.E. Schevill. 1982. Observations of right whales (*Eubalaena glacialis*) in Cape Cod waters. Fish. Bull. 80(4): 875-880.
- Watkins, W.A., K.E. Moore, J. Sigurjonsson, D. Wartzok, and G. Notarbartolo di Sciara. 1984. Fin whale (*Balaenoptera physalus*) tracked by radio in the Irminger Sea. Rit Fiskideildar 8(1): 1-14.
- Watling, L. 1998. Benthic fauna of soft substrates in the Gulf of Maine. Pp. 20-29 in: Effects of fishing gear on the sea floor of New England, E.M. Dorsey and J. Pederson (eds.). MIT Sea Grant Pub. 98-4.
- Watling, L., R.H. Findlay, L.M. Mayer, and D.F. Schick. 2001. Impact of a scallop drag on the sediment chemistry, microbiota, and faunal assemblages of a shallow subtidal marine benthic community. Journal of Sea Research 46:309-324.
- West, T.L., W.G. Ambrose, Jr., and G.A. Skilleter. 1994. A review of the effects of fish harvesting practices on the benthos and bycatch: implications and recommendations for North Carolina. Albemarle-Pamlico Estuarine Study, Raleigh, NC, U.S. Environmental Protection Agency and NC Department of Health, Environment and Natural Resources. Report No. 94-06.
- Wiebe, P.H., E.H. Backus, R.H. Backus, D.A. Caron, P.M. Glibert, J.F. Grassle, K. Powers, and J.B. Waterbury. 1987. Biological oceanography. Pp. 140-201 in: J.D. Milliman and W.R. Wright (eds), The marine environment of the U.S. Atlantic continental slope and rise. Jones and Bartlett Publishers, Inc., Boston, MA.
- Wigley, R.L. and R.B. Theroux. 1981. Atlantic continental shelf and slope of the United States – macrobenthic invertebrate fauna of the middle Atlantic bight region – faunal composition and quantitative distribution. Geological Survey Professional Paper 529-N, United States Department of the Interior. 198 p.

- Wigley, Susan E. 1996. The Lorenz curve method applied to NEFSC bottom trawl survey data, A report of Northeast Regional Stock Assessment Workshop No. 21. Northeast Fisheries Science Center, Woods Hole, MA. Center Reference Document CRD 96-05f.
- Wiley, D.N., R.A. Asmutis, T.D. Pitchford, and D.P. Gannon. 1995. Stranding and mortality of humpback whales, *Megaptera novaengliae*, in the mid-Atlantic and southeast United States, 1985-1992. Fish. Bull., U.S. 93:196-205.
- Wilhelm, Kurt A. 2001. An analysis of selected trips in the Gulf of Maine cod fishery, January 2000 through January 2001. Unpublished manuscript. National Marine Fisheries Service, Northeast Regional Office. Gloucester, MA.
- Winn, H.E., C.A. Price, and P.W. Sorensen. 1986. The distributional biology of the right whale (*Eubalaena glacialis*) in the western North Atlantic. Rep. Int. Whal. Comm.. Spec. Iss. 10:129-138.
- Witman, J.D. 1998. Natural disturbance and colonization on subtidal hard substrates in the Gulf of Maine. In E.M. Dorsey and J. Pederson, eds. Effects of Fishing Gear on the Sea Floor of New England. Boston, MA: Conservation Law Foundation. 30-37.
- Woods Hole Oceanographic Institution, 2001. Second Symposium on Fisheries, Oceanography and Society, Marine Protected Areas: Design and implementation for Conservation and Fisheries Restoration, 68p.
- Worthington, L.V. 1976. On the North Atlantic circulation. Johns Hopkins Oceanographic Studies No. 6. The Johns Hopkins University Press, Baltimore, MD. 110 p.
- Wright, W.R. and L.V. Worthington. 1970. The water masses of the North Atlantic Ocean: a volumetric census of temperature and salinity. Serial Atlas of the Marine Environment. American Geological Society Folio No. 19.
- Wynne, K. and M. Schwartz. 1999. Guide to marine mammals and turtles of the U.S. Atlantic and Gulf of Mexico. Rhode Island Sea Grant, Narragansett. 115pp.
- Yeung, C. 1999. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. U.S. Dep. Commer. NOAA Tech. Memo. NMFS-NEFSC-430, 26pp.
- Zug, G. R. and J.F. Parham. 1996. Age and growth in leatherback turtles, *Dermochelys coriacea*: a skeletochronological analysis. Chelonian Conservation and Biology. 2(2): 244-249.

8.2 Glossary

Adult stage: One of several marked phases or periods in the development and growth of many animals. In vertebrates, the life history stage where the animal is capable of reproducing, as opposed to the juvenile stage.

Adverse effect: Any impact that reduces quality and/or quantity of EFH. May include direct or indirect physical, chemical, or biological alterations of the waters or substrate and loss of, or injury to, benthic organisms, prey species and their habitat, and other ecosystem components, if such modifications reduce the quality and or quantity of EFH. Adverse effects to EFH may result from actions occurring within EFH or outside of EFH and may include sites-specific or habitat wide impacts, including individual, cumulative, or synergistic consequences of actions.

Aggregation: A group of animals or plants occurring together in a particular location or region.

Anadromous species: fish that spawn in fresh or estuarine waters and migrate to ocean waters

Amphipods: A small crustacean of the order Amphipoda, such as the beach flea, having a laterally compressed body with no carapace.

Anaerobic sediment: Sediment characterized by the absence of free oxygen.

Anemones: Any of numerous flowerlike marine coelenterates of the class Anthozoa, having a flexible cylindrical body and tentacles surrounding a central mouth.

Annual total mortality: Rate of death expressed as the fraction of a cohort dying over a period compared to the number alive at the beginning of the period ($\#$ total deaths during year / numbers alive at the beginning of the year). Optimists convert death rates into annual survival rate using the relationship $S=1-A$.

ASPIC (A Surplus Production Model Incorporating Covariates): A non-equilibrium surplus production model developed by Prager (1995). ASPIC was frequently used by the Overfishing Definition Panel to define B_{MSY} and F_{MSY} reference points. The model output was also used to estimate rebuilding timeframes for the Amendment 9 control rules.

Bay: An inlet of the sea or other body of water usually smaller than a gulf; a small body of water set off from the main body; e.g. Ipswich Bay in the Gulf of Maine.

Benthic community: *Benthic* means the bottom habitat of the ocean, and can mean anything as shallow as a salt marsh or the intertidal zone, to areas of the bottom that are several miles deep in the ocean. *Benthic community* refers to those organisms that live in and on the bottom. (*In* meaning they live within the substrate; e.g. within the sand or mud found on the bottom. See *Benthic infauna*, below)

Benthic infauna: See *Benthic community*, above. Those organisms that live *in* the bottom sediments (sand, mud, gravel, etc.) of the ocean. As opposed to *benthic epifauna*, that live *on* the surface of the bottom sediments.

Benthivore: Usually refers to fish that feed on benthic or bottom dwelling organisms.

Berm: A narrow ledge typically at the top or bottom of a slope; e.g. a berm paralleling the shoreline caused by wave action on a sloping beach; also an elongated mound or wall of earth.

Biogenic habitats: Ocean habitats whose physical structure is created or produced by the animals themselves; e.g., coral reefs.

Biomass: The total mass of living matter in a given unit area or the weight of a fish stock or portion thereof. Biomass can be listed for beginning of year (Jan-1), Mid-Year, or mean (average during the entire year). In addition, biomass can be listed by age group (numbers at age * average weight at age) or summarized by groupings (e.g., age 1+, ages 4+ 5, etc). See also spawning stock biomass, exploitable biomass, and mean biomass.

B_{MSY}: The stock biomass that would produce MSY when fished at a fishing mortality rate equal to F_{MSY}. For most stocks, B_{MSY} is about ½ of the carrying capacity. The proposed overfishing definition control rules call for action when biomass is below ¼ or ½ B_{MSY}, depending on the species.

B_{threshold}: 1) A limit reference point for biomass that defines an unacceptably low biomass i.e., puts a stock at high risk (recruitment failure, depensation, collapse, reduced long term yields, etc). 2) A biomass threshold that the SFA requires for defining when a stock is overfished. A stock is overfished if its biomass is below B_{threshold}. A determination of overfished triggers the SFA requirement for a rebuilding plan to achieve B_{target} as soon as possible, usually not to exceed 10 years except certain requirements are met. In Amendment 9 control rules, B_{threshold} is often defined as either 1/2B_{MSY} or 1/4 B_{MSY}. B_{threshold} is also known as B_{minimum}.

B_{target}: A desirable biomass to maintain fishery stocks. This is usually synonymous with B_{MSY} or its proxy.

Biomass weighted F: A measure of fishing mortality that is defined as an average of fishing mortality at age weighted by biomass at age for a ranges of ages within the stock (e.g., ages 1+ biomass weighted F is a weighted average of the mortality for ages 1 and older, age 3+ biomass weighted is a weighted average for ages 3 and older). Biomass weighted F can also be calculated using catch in weight over mean biomass. See also fully-recruited F.

Biota: All the plant and animal life of a particular region.

Bivalve: A class of mollusks having a soft body with platelike gills enclosed within two shells hinged together; e.g., clams, mussels.

Bottom roughness: The inequalities, ridges, or projections on the surface of the seabed that are caused by the presence of bedforms, sedimentary structures, sedimentary particles, excavations, attached and unattached organisms, or other objects; generally small scale features.

Bottom tending mobile gear: All fishing gear that operates on or near the ocean bottom that is actively worked in order to capture fish or other marine species. Some examples of bottom tending mobile gear are otter trawls and dredges.

Bottom tending static gear: All fishing gear that operates on or near the ocean bottom that is not actively worked; instead, the effectiveness of this gear depends on species moving to the gear which is set in a particular manner by a vessel, and later retrieved. Some examples of bottom tending static gear are gillnets, traps, and pots.

Boulder reef: An elongated feature (a chain) of rocks (generally piled boulders) on the seabed.

Bryozoans: Phylum aquatic organisms, living for the most part in colonies of interconnected individuals. A few to many millions of these individuals may form one colony. Some bryozoans encrust rocky surfaces, shells, or algae others form lacy or fan-like colonies that in some regions may form an abundant

component of limestones. Bryozoan colonies range from millimeters to meters in size, but the individuals that make up the colonies are rarely larger than a millimeter. Colonies may be mistaken for hydroids, corals or seaweed.

Burrow: A hole or excavation in the sea floor made by an animal (as a crab, lobster, fish, burrowing anemone) for shelter and habitation.

Bycatch: (v.) the capture of nontarget species in directed fisheries which occurs because fishing gear and methods are not selective enough to catch only target species; (n.) fish which are harvested in a fishery but are not sold or kept for personal use, including economic discards and regulatory discards but not fish released alive under a recreational catch and release fishery management program.

Capacity: the level of output a fishing fleet is able to produce given specified conditions and constraints. Maximum fishing capacity results when all fishing capital is applied over the maximum amount of available (or permitted) fishing time, assuming that all variable inputs are utilized efficiently.

Catch: The sum total of fish killed in a fishery in a given period. Catch is given in either weight or number of fish and may include landings, unreported landings, discards, and incidental deaths.

Coarse sediment: Sediment generally of the sand and gravel classes; not sediment composed primarily of mud; but the meaning depends on the context, e.g. within the mud class, silt is coarser than clay.

Commensalism: See *Mutualism*. An interactive association of two species where one benefits in some way, while the other species is in no way affected by the association.

Continental shelf waters: The waters overlying the continental shelf, which extends seaward from the shoreline and deepens gradually to the point where the sea floor begins a slightly steeper descent to the deep ocean floor; the depth of the shelf edge varies, but is approximately 200 meters in many regions.

Control rule: A pre-determined method for determining fishing mortality rates based on the relationship of current stock biomass to a biomass target. Amendment 9 overfishing control rules define a target biomass (B_{MSY} or proxy) as a management objective. The biomass threshold ($B_{threshold}$ or B_{min}) defines a minimum biomass below which a stock is considered overfished.

Cohort: see yearclass.

Crustaceans: Invertebrates characterized by a hard outer shell and jointed appendages and bodies. They usually live in water and breathe through gills. Higher forms of this class include lobsters, shrimp and crawfish; lower forms include barnacles.

Diatoms: Small mobile plants (algæ) with silicified (silica, sand, quartz) skeletons. They are among the most abundant phytoplankton in cold waters, and an important part of the food chain.

Days absent: an estimate by port agents of trip length. This data was collected as part of the NMFS weighout system prior to May 1, 1994.

Days-at-sea (DAS): the total days, including steaming time that a boat spends at sea to fish.

Demersal species: Most often refers to fish that live on or near the ocean bottom. They are often called benthic fish, groundfish, or bottom fish.

Discards: animals returned to sea after being caught; see Bycatch (n.)

Dissolved nutrients: Non-solid nutrients found in a liquid.

Echinoderms: A member of the Phylum Echinodermata. Marine animals usually characterised by a five-fold symmetry, and possessing an internal skeleton of calcite plates, and a complex water vascular system. Includes echinoids (sea urchins), crinoids (sea lillies) and asteroids (starfish).

Ecosystem-based management: a management approach that takes major ecosystem components and services—both structural and functional—into account, often with a multispecies or habitat perspective

Egg stage: One of several marked phases or periods in the development and growth of many animals. The life history stage of an animal that occurs after reproduction and refers to the developing embryo, its food store, and sometimes jelly or albumen, all surrounded by an outer shell or membrane. Occurs before the *larval* or *juvenile stage*.

Elasmobranch: Any of numerous fishes of the class Chondrichthyes characterized by a cartilaginous skeleton and placoid scales: sharks; rays; skates.

Embayment: A bay or an indentation in a coastline resembling a bay.

Emergent epifauna: See *Epifauna*. Animals living upon the bottom that extend a certain distance above the surface.

Epifauna: See *Benthic infauna*. *Epifauna* are animals that live on the surface of the substrate, and are often associated with surface structures such as rocks, shells, vegetation, or colonies of other animals.

Essential Fish Habitat (EFH): Those waters and substrate necessary to fish for spawning, breeding, feeding, or growth to maturity. The EFH designation for most managed species in this region is based on a legal text definition and geographical area that are described in the Habitat Omnibus Amendment (1998).

Estuarine area: The area of an estuary and its margins; an area characterized by environments resulting from the mixing of river and sea water.

Estuary: A water passage where the tide meets a river current; especially an arm of the sea at the lower end of a river; characterized by an environment where the mixing of river and seawater causes marked variations in salinity and temperature in a relatively small area.

Eutrophication: A set of physical, chemical, and biological changes brought about when excessive nutrients are released into the water.

Euphotic zone: The zone in the water column where at least 1% of the incident light at the surface penetrates.

Exclusive Economic Zone (EEZ): a zone in which the inner boundary is a line coterminous with the seaward boundary of each of the coastal States and the outer boundary is line 200 miles away and parallel to the inner boundary

Exempt fisheries: Any fishery determined by the Regional Director to have less than 5 percent regulated species as a bycatch (by weight) of total catch according to 50 CFR ?648.80(a)(7).

Exploitable biomass: The biomass of fish in the portion of the population that is vulnerable to fishing.

Exploitation pattern: Describes the fishing mortality at age as a proportion of fully recruited F (full vulnerability to the fishery). Ages that are fully vulnerable experience 100% of the fully recruited F and are

termed fully recruited. Ages that are only partially vulnerable experience a fraction of the fully recruited F and are termed partially recruited. Ages that are not vulnerable to the fishery (including discards) experience no mortality and are considered pre-recruits. Also known as the partial recruitment pattern, partial recruitment vector or fishery selectivity.

Exploitation rate (u): The fraction of fish in the exploitable population killed during the year by fishing. This is an annual rate compared to F , which is an instantaneous rate. For example, if a population has 1,000,000 fish large enough to be caught and 550,000 are caught (landed and discarded) then the exploitation rate is 55%.

Fathom: A measure of length, containing six feet; the space to which a man can extend his arms; used chiefly in measuring cables, cordage, and the depth of navigable water by soundings.

Fishing mortality (F): A measurement of the rate of removal of fish from a population caused by fishing. This is usually expressed as an instantaneous rate (F) and is the rate at which fish are harvested at any given point in a year. Instantaneous fishing mortality rates can be either fully recruited or biomass weighted. Fishing mortality can also be expressed as an exploitation rate (see exploitation rate) or less commonly, as a conditional rate of fishing mortality (m , fraction of fish removed during the year if no other competing sources of mortality occurred. Lower case m should not be confused with upper case M , the instantaneous rate of natural mortality).

$F_{0.1}$: a conservative fishing mortality rate calculated as the F associated with 10 percent of the slope at origin of the yield-per-recruit curve.

F_{MAX} : a fishing mortality rate that maximizes yield per recruit. F_{MAX} is less conservative than $F_{0.1}$.

F_{MSY} : a fishing mortality rate that would produce MSY when the stock biomass is sufficient for producing MSY on a continuing basis.

$F_{threshold}$: 1) The maximum fishing mortality rate allowed on a stock and used to define overfishing for status determination. Amendment 9 frequently uses F_{MSY} or F_{MSY} proxy for $F_{threshold}$. 2) The maximum fishing mortality rate allowed for a given biomass as defined by a control rule.

Fishing effort: the amount of time and fishing power used to harvest fish. Fishing power is a function of gear size, boat size and horsepower.

Fluidized: When a normally non-fluid material is made to act like a fluid; e.g. gravity, shaking or injection of a fluid.

Framework adjustments: adjustments within a range of measures previously specified in a fishery management plan (FMP). A change usually can be made more quickly and easily by a framework adjustment than through an amendment. For plans developed by the New England Council, the procedure requires at least two Council meetings including at least one public hearing and an evaluation of environmental impacts not already analyzed as part of the FMP.

Furrow: A trench in the earth made by a plow; something that resembles the track of a plow, as a marked narrow depression; a groove with raised edges.

Glacial moraine: A sedimentary feature deposited from glacial ice; characteristically composed of unsorted clay, sand, and gravel. Moraines typically are hummocky or ridge-shaped and are located along the sides and at the fronts of glaciers.

Glacial till: Unsorted sediment (clay, sand, and gravel mixtures) deposited from glacial ice.

Grain size: the size of individual sediment particles that form a sediment deposit; particles are separated into size classes (e.g. very fine sand, fine sand, medium sand, among others); the classes are combined into broader categories of mud, sand, and gravel; a sediment deposit can be composed of few to many different grain sizes.

Growth overfishing: Fishing at an exploitation rate or at an age at entry that reduces potential yields from a cohort but does not reduce reproductive output (see recruitment overfishing).

Halocline: The zone of the ocean in which salinity increases rapidly with depth.

Habitat complexity: Describes or measures a habitat in terms of the variability of its characteristics and its functions, which can be biological, geological, or physical in nature. Refers to how complex the physical structure of the habitat is. A bottom habitat with *structure-forming organisms*, along with other three dimensional objects such as boulders, is more complex than a flat, featureless, bottom.

Highly migratory species: tuna species, marlin, oceanic sharks, sailfishes, and swordfish

Hydroids: Generally, animals of the Phylum Cnidaria, Class Hydrozoa; most hydroids are bush-like polyps growing on the bottom and feed on plankton, they reproduce asexually and sexually.

Immobile epifaunal species: See *epifauna*. Animals living on the surface of the bottom substrate that, for the most part, remain in one place.

Individual Fishing Quota (IFQ): federal permit under a limited access system to harvest a quantity of fish, expressed by a unit or units representing a percentage of the total allowable catch of a fishery that may be received or held for exclusive use by an individual person or entity

Juvenile stage: One of several marked phases or periods in the development and growth of many animals. The life history stage of an animal that comes between the *egg* or *larval stage* and the *adult stage*; juveniles are considered immature in the sense that they are not yet capable of reproducing, yet they differ from the larval stage because they look like smaller versions of the adults.

Landings: The portion of the catch that is harvested for personal use or sold.

Land runoff: The part of precipitation, snowmelt, or irrigation water that reaches streams (and thence the sea) by flowing over the ground, or the portion of rain or snow that does not percolate into the ground and is discharged into streams instead.

Larvae stage: One of several marked phases or periods in the development and growth of many animals. The first stage of development after hatching from the *egg* for many fish and invertebrates. This life stage looks fundamentally different than the juvenile and adult stages, and is incapable of reproduction; it must undergo metamorphosis into the juvenile or adult shape or form.

Lethrinids: Fish of the genus *Lethrinus*, commonly called emperors or nor'west snapper, are found mainly in Australia's northern tropical waters. Distinctive features of Lethrinids include thick lips, robust canine teeth at the front of the jaws, molar-like teeth at the side of the jaws and cheeks without scales. Lethrinids are carnivorous bottom-feeding fish with large, strong jaws.

Limited-access permits: permits issued to vessels that met certain qualification criteria by a specified date (the "control date").

Lutjanids: Fish of the genus of the Lutjanidae: snappers. Marine; rarely estuarine. Some species do enter freshwater for feeding. Tropical and subtropical: Atlantic, Indian and Pacific Oceans.

Macrobenthos: See *Benthic community* and *Benthic infauna*. Benthic organisms whose shortest dimension is greater than or equal to 0.5 mm.

Maturity ogive: A mathematical model used to describe the proportion mature at age for the entire population. A_{50} is the age where 50% of the fish are mature.

Mean biomass: The average number of fish within an age group alive during a year multiplied by average weight at age of that age group. The average number of fish during the year is a function of starting stock size and mortality rate occurring during the year. Mean biomass can be aggregated over several ages to describe mean biomass for the stock. For example the mean biomass summed for ages 1 and over is the 1^+ mean biomass; mean biomass summed across ages 3 and over is 3^+ mean biomass.

Megafaunal species: The component of the fauna of a region that comprises the larger animals, sometimes defined as those weighing more than 100 pounds.

Mesh selectivity ogive: A mathematical model used to describe the selectivity of a mesh size (proportion of fish at a specific length retained by mesh) for the entire population. L_{25} is the length where 25% of the fish encountered are retained by the mesh. L_{50} is the length where 50% of the fish encountered are retained by the mesh.

Meter: A measure of length, equal to 39.37 English inches, the standard of linear measure in the metric system of weights and measures. It was intended to be, and is very nearly, the ten millionth part of the distance from the equator to the north pole, as ascertained by actual measurement of an arc of a meridian.

Metric ton: A unit of weight equal to a thousand kilograms (1kgs = 2.2 lbs.). A metric ton is equivalent to 2,205 lbs. A thousand metric tons is equivalent to 2.2 million lbs.

Microalgal: Small microscopic types of algae such as the green algae.

Microbial: Microbial means of or relating to microorganisms.

Minimum spawning stock threshold: the minimum spawning stock size (or biomass) below which there is a significantly lower chance that the stock will produce enough new fish to sustain itself over the long term.

Mobile organisms: organisms that are not confined or attached to one area or place, that can move on their own, are capable of movement, or are moved (often passively) by the action of the physical environment (waves, currents, etc.).

Molluscs: Common term for animals of the phylum Mollusca. Includes groups such as the bivalves (mussels, oysters etc.), cephalopods (squid, octopus etc.) and gastropods (abalone, snails). Over 80,000 species in total with fossils back to the Cambrian period.

Mortality: see Annual total mortality (A), Exploitation rate (u), Fishing mortality (F), Natural mortality (M), and instantaneous total mortality (Z).

Motile: Capable of self-propelled movement. A term that is sometimes used to distinguish between certain types of organisms found in water.

Multispecies: the group of species managed under the Northeast Multispecies Fishery Management Plan. This group includes whiting, red hake and ocean pout plus the regulated species (cod, haddock, pollock,

yellowtail flounder, winter flounder, witch flounder, American plaice, windowpane flounder, white hake and redfish).

Mutualism: See *Commensalism*. A symbiotic interaction between two species in which both derive some benefit.

Natural disturbance: A change caused by natural processes; e.g. in the case of the seabed, changes can be caused by the removal or deposition of sediment by currents; such natural processes can be common or rare at a particular site.

Natural mortality: A measurement of the rate of death from all causes other than fishing such as predation, disease, starvation, and pollution. Commonly expressed as an instantaneous rate (M). The rate of natural mortality varies from species to species, but is assumed to be $M=0.2$ for the five critical stocks. The natural mortality rate can also be expressed as a conditional rate (termed n and not additive with competing sources of mortality such as fishing) or as annual expectation of natural death (termed v and additive with other annual expectations of death).

Nearshore area: The area extending outward an indefinite but usually short distance from shore; an area commonly affected by tides and tidal and storm currents, and shoreline processes.

Nematodes: a group of elongated, cylindrical worms belonging to the phylum Nematodea, also called thread-worms or eel-worms. Some non-marine species attack roots or leaves of plants, others are parasites on animals or insects.

Nemertean: Proboscis worms belonging to the phylum Nemertea, and are soft unsegmented marine worms that have a threadlike proboscis and the ability to stretch and contract.

Nemipterids: Fishes of the Family Nemipteridae, the threadfin breams or whiptail breams. Distribution: Tropical and sub-tropical Indo-West Pacific.

Northeast Shelf Ecosystem: The Northeast U.S. Shelf Ecosystem has been described as including the area from the Gulf of Maine south to Cape Hatteras, extending from the coast seaward to the edge of the continental shelf, including the slope sea offshore to the Gulf Stream.

Northwest Atlantic Analysis Area (NAAA): A spatial area developed for analysis purposes only. The boundaries of this the area are within the 500 fathom line to the east, the coastline to the west, the Hague line to the north, and the North Carolina/ South Carolina border to the south. The area is approximately 83,550 square nautical miles, and is used as the denominator in the EFH analysis to determine the percent of sediment, EFH, and biomass contained in an area, as compared to the total NAAA.

Nutrient budgets: An accounting of nutrient inputs to and production by a defined ecosystem (e.g., salt marsh, estuary) versus utilization within and export from the ecosystem.

Observer: any person required or authorized to be carried on a vessel for conservation and management purposes by regulations or permits under this Act

Oligochaetes: See *Polychaetes*. Oligochaetes are worms in the phylum Annelida having bristles borne singly along the length of the body.

Open access: describes a fishery or permit for which there is no qualification criteria to participate. Open-access permits may be issued with restrictions on fishing (for example, the type of gear that may be used or the amount of fish that may be caught).

Opportunistic species: Species that colonize disturbed or polluted sediments. These species are often small, grow rapidly, have short life spans, and produce many offspring.

Optimum Yield (OY): the amount of fish which A) will provide the greatest overall benefit to the nation, particularly with respect to food production and recreational opportunities, and taking into account the protection of marine ecosystems; B) is prescribed as such on the basis of the maximum sustainable yield from the fishery, as reduced by any relevant economic, social, or ecological factor; and C) in the case of an overfished fishery, provides for rebuilding to a level consistent with producing the maximum sustainable yield in such fishery

Organic matter: Material of, relating to, or derived from living organisms.

Overfished: A condition defined when stock biomass is below minimum biomass threshold and the probability of successful spawning production is low.

Overfishing: A level or rate of fishing mortality that jeopardizes the long-term capacity of a stock or stock complex to produce MSY on a continuing basis.

Peat bank: A bank feature composed of partially carbonized, decomposed vegetable tissue formed by partial decomposition of various plants in water; may occur along shorelines.

Pelagic gear: Mobile or static fishing gear that is not fixed, and is used within the water column, not on the ocean bottom. Some examples are mid-water trawls and pelagic longlines.

Phytoplankton: Microscopic marine plants (mostly algae and diatoms) which are responsible for most of the photosynthetic activity in the oceans.

Piscivore: A species feeding preferably on fish.

Planktivore: An animal that feeds on plankton.

Polychaetes: Polychaetes are segmented worms in the phylum Annelida. Polychaetes (poly-chaetae = many-setae) differ from other annelids in having many setae (small bristles held in tight bundles) on each segment.

Porosity: The amount of free space in a volume of a material; e.g. the space that is filled by water between sediment particles in a cubic centimeter of seabed sediment.

Possession-limit-only permit: an open-access permit (see above) that restricts the amount of multispecies a vessel may retain (currently 500 pounds of "regulated species").

Pre-recruits: Fish in size or age groups that are not vulnerable to the fishery (including discards).

Prey availability: The availability or accessibility of prey (food) to a predator. Important for growth and survival.

Primary production: The synthesis of organic materials from inorganic substances by photosynthesis.

Recovery time: The period of time required for something (e.g. a habitat) to achieve its former state after being disturbed.

Recruitment: the amount of fish added to the fishery each year due to growth and/or migration into the fishing area. For example, the number of fish that grow to become vulnerable to fishing gear in one year

would be the recruitment to the fishery. “Recruitment” also refers to new year classes entering the population (prior to recruiting to the fishery).

Recruitment overfishing: fishing at an exploitation rate that reduces the population biomass to a point where recruitment is substantially reduced.

Regulated groundfish species: cod, haddock, pollock, yellowtail flounder, winter flounder, witch flounder, American plaice, windowpane flounder, white hake and redfish. These species are usually targeted with large-mesh net gear.

Relative exploitation: an index of exploitation derived by dividing landings by trawl survey biomass. This measure does not provide an absolute magnitude of exploitation but allows for general statements about trends in exploitation.

Retrospective pattern: A pattern of systematic over-estimation or underestimation of terminal year estimates of stock size, biomass or fishing mortality compared to that estimate for that same year when it occurs in pre-terminal years.

Riverine area: The area of a river and its banks.

Saurids: Fish of the family Scomberesocidae, the sauries or needlefishes. Distribution: tropical and temperate waters.

Scavenging species: An animal that consumes dead organic material.

Sea whips: A coral that forms long flexible structures with few or no branches and is common on Atlantic reefs.

Sea pens: An animal related to corals and sea anemones with a featherlike form.

Sediment: Material deposited by water, wind, or glaciers.

Sediment suspension: The process by which sediments are suspended in water as a result of disturbance.

Sedentary: See *Motile* and *Mobile organisms*. Not moving. Organisms that spend the majority of their lives in one place.

Sedimentary bedforms: Wave-like structures of sediment characterized by crests and troughs that are formed on the seabed or land surface by the erosion, transport, and deposition of particles by water and wind currents; e.g. ripples, dunes.

Sedimentary structures: Structures of sediment formed on the seabed or land surface by the erosion, transport, and deposition of particles by water and wind currents; e.g. ripples, dunes, buildups around boulders, among others.

Sediment types: Major combinations of sediment grain sizes that form a sediment deposit, e.g. mud, sand, gravel, sandy gravel, muddy sand, among others.

Spawning adult stage: See *adult stage*. Adults that are currently producing or depositing eggs.

Spawning stock biomass (SSB): the total weight of fish in a stock that sexually mature, i.e., are old enough to reproduce.

Species assemblage: Several species occurring together in a particular location or region

Species composition: A term relating the relative abundance of one species to another using a common measurement; the proportion (percentage) of various species in relation to the total on a given area.

Species diversity: The number of different species in an area and their relative abundance

Species richness: See *Species diversity*. A measurement or expression of the number of species present in an area; the more species present, the higher the degree of species richness.

Species with vulnerable EFH: If a species was determined to be “highly” or “moderately” vulnerable to bottom tending gears (otter trawls, scallop dredges, or clam dredges) then it was included in the list of species with vulnerable EFH. Currently there are 23 species and life stages that are considered to have vulnerable EFH for this analysis.

Status Determination: A determination of stock status relative to $B_{\text{threshold}}$ (defines overfished) and $F_{\text{threshold}}$ (defines overfishing). A determination of either overfished or overfishing triggers a SFA requirement for rebuilding plan (overfished), ending overfishing (overfishing) or both.

Stock: A grouping of fish usually based on genetic relationship, geographic distribution and movement patterns. A region may have more than one stock of a species (for example, Gulf of Maine cod and Georges Bank cod). A species, subspecies, geographical grouping, or other category of fish capable of management as a unit.

Stock assessment: determining the number (abundance/biomass) and status (life-history characteristics, including age distribution, natural mortality rate, age at maturity, fecundity as a function of age) of individuals in a stock

Structure-forming organisms: Organisms, such as corals, colonial bryozoans, hydroids, sponges, mussel beds, oyster beds, and seagrass that by their presence create a three-dimensional physical structure on the bottom. See *biogenic habitats*.

Submerged aquatic vegetation: Rooted aquatic vegetation, such as seagrasses, that cannot withstand excessive drying and therefore live with their leaves at or below the water surface in shallow areas of estuaries where light can penetrate to the bottom sediments. SAV provides an important habitat for young fish and other aquatic organisms.

Surficial sediment: Sediment forming the sea floor or land surface; thickness of the surficial layer may vary.

Surplus production: Production of new stock biomass defined by recruitment plus somatic growth minus biomass loss due to natural deaths. The rate of surplus production is directly proportional to stock biomass and its relative distance from the maximum stock size at carrying capacity (K). B_{MSY} is often defined as the biomass that maximizes surplus production rate.

Surplus production models: A family of analytical models used to describe stock dynamics based on catch in weight and CPUE time series (fishery dependent or survey) to construct stock biomass history. These models do not require catch at age information. Model outputs may include stock biomass history, biomass weighted fishing mortality rates, MSY , F_{MSY} , B_{MSY} , K , (maximum population biomass where stock growth and natural deaths are balanced) and r (intrinsic rate of increase).

Survival rate (S): Rate of survival expressed as the fraction of a cohort surviving the a period compared to number alive at the beginning of the period (# survivors at the end of the year / numbers alive at the

beginning of the year). Pessimists convert survival rates into annual total mortality rate using the relationship $A=1-S$.

Survival ratio (R/SSB): an index of the survivability from egg to age-of-recruitment. Declining ratios suggest that the survival rate from egg to age-of-recruitment is declining.

TAC: Total allowable catch. This value is calculated by applying a target fishing mortality rate to exploitable biomass.

Taxa: The plural of taxon. Taxon is a named group of organisms of any rank, such as a particular species, family, or class.

Ten-minute- “squares” of latitude and longitude (TMS): Are a measure of geographic space. The actual size of a ten-minute-square varies depending on where it is on the surface of the earth, but in general each square is approximately 70-80 square nautical miles in this region. This is the spatial area that EFH designations, biomass data, and some of the effort data have been binned into for analysis purposes in various sections of this document.

Topography: The depiction of the shape and elevation of land and sea floor surfaces.

Total mortality: The rate of mortality from all sources (fishing, natural, pollution) Total mortality can be expressed as an instantaneous rate (called Z and equal to $F + M$) or Annual rate (called A and calculated as the ratio of total deaths in a year divided by number alive at the beginning of the year)

Trophic guild: Trophic is defined as the feeding level within a system that an organism occupies; e.g., predator, herbivore. A guild is defined as a group of species that exploit the same class of environmental resources in a similar way. The trophic guild is a utilitarian concept covering both structure and organization that exists between the structural categories of trophic groups and species.

Turbidity: Relative water clarity; a measurement of the extent to which light passing through water is reduced due to suspended materials.

Two-bin (displacement) model: a model used to estimate the effects of area closures. This model assumes that effort from the closed areas (first bin) is displaced to the open areas (second bin). The total effort in the system is then applied to the landings-per-unit-effort (LPUE) in open areas to obtain a projected catch. The percent reduction in catch is calculated as a net result.

Vulnerability: In order to evaluate the potential adverse effects of fishing on EFH, the vulnerability of each species EFH was determined. This analysis defines vulnerability as the likelihood that the functional value of EFH would be adversely affected as a result of fishing with different gear types. A number of criteria were considered in the evaluation of the vulnerability of EFH for each life stage including factors like the function of habitat for shelter, food and/or reproduction.

Yield-per-recruit (YPR): the expected yield (weight) of individual fish calculated for a given fishing mortality rate and exploitation pattern and incorporating the growth characteristics and natural mortality.

Yearclass: also called cohort. Fish that were spawned in the same year. By convention, the “birth date” is set to January 1st and a fish must experience a summer before turning 1. For example, winter flounder that were spawned in February-April 1997 are all part of the 1997 cohort (or year-class). They would be considered age 0 in 1997, age 1 in 1998, etc. A summer flounder spawned in October 1997 would have its birth date set to the following January 1 and would be considered age 0 in 1998, age 1 in 1999, etc.

Z: instantaneous rate of total mortality. The components of Z are additive (i.e., $Z = F+M$)

Zooplankton: See *Phytoplankton*. Small, often microscopic animals that drift in currents. They feed on detritus, phytoplankton, and other zooplankton. They are preyed upon by fish, shellfish, whales, and other zooplankton.

8.3 Table of Acronyms

The table of acronyms can be found after the table of contents.

8.4 List of Public Meetings

Amendment 13 issues were discussed at the following meetings.

	Meeting Type	Meeting Location	Meeting Dates
2000	Council Meeting	Tavern on Harbor, Gloucester	3/22/2000
	Council Meeting	Providence Biltmore, PVD, RI	5/3 & 5/4/2000
	Council Meeting	Sheraton, Portsmouth, NH	6/14 & 6/15/2000
	Council Meeting	DoubleTree Hotel, Portland, ME	7/26 & 7/27/2000
	Council Meeting	Seaport Inn, Fairhaven, MA	9/25 - 9/27/2000
	Council Meeting	Tavern on Harbor, Gloucester	11/6 - 11/8/2000
	Amendment 13 Info Mtg	Gloucester Seafood Display	4/27/2000
	Amendment 13 Info Mtg	Mass Maritime, Buzzards Bay	5/11/2000
	Amendment 13 Info Mtg	Rockland Middle School, ME	5/18/2000
	Amendment 13 Info Mtg	Radisson Hotel, New London, CT	5/25/2000
	Amendment 13 Info Mtg	Fish Family Asst Ctr, New Bedford	6/1/2000
	Amendment 13 Info Mtg	Gloucester, MA	6/2/2000
	Oversight Committee	Holiday Inn, Peabody, MA	1/14/2000
	Oversight Committee	Yokens Ctr, Portsmouth, NH	3/15/2000
	Oversight Committee	Holiday Inn, Mansfield, MA	4/11/2000
	Oversight Committee/Advisory Panel	Sheraton Ferncroft, Danvers	2/16/2000
	Oversight Committee/Advisory Panel	Sheraton Colonial, Wakefield, MA	5/1 & 5/2/2000
	Oversight Committee/Advisory Panel	York Harbor Inn, York, ME	5/9/2000
	Oversight Committee/Advisory Panel	Holiday Inn, Peabody, MA	5/22/2000
	Oversight Committee/Advisory Panel	Howard Johnson's, Portsmouth, NH	6/13/2000
	Oversight Committee/Advisory Panel	Radisson Hotel, New London, CT	6/20/2000
	Oversight Committee/Advisory Panel	Trade Winds, Rockland, ME	7/6/2000
	Oversight Committee/Advisory Panel	Sheraton Ferncroft, Danvers	7/13/2000
	Oversight Committee/Advisory Panel	Seaport Inn, Fairhaven, MA	7/19/2000
	Advisory Panel	Holiday Inn, Peabody, MA	1/13/2000
	Advisory Panel	Sheraton Airport, Warwick, RI	6/27/2000
	PDT	NEFMC, Newburyport, MA	8/3/2000
	PDT	NEFSC, Woods Hole, MA	9/25/2000
	PDT	NEFSC, Woods Hole, MA	10/18/2000
	PDT	Aquarium Library, Woods Hole, MA	12/7/2000
	PDT	NEFSC, Woods Hole, MA	12/14/2000
	Social Impact Informational Meeting (Public Hearing)	Seafood Display Auction, Gloucester	11/1/2000
	Social Impact Informational Meeting (Public Hearing)	Eldridge Library, Chatham, MA	11/2/2000
	Social Impact Informational Meeting (Public Hearing)	Portsmouth, NH	11/6/2000
	Social Impact Informational Meeting (Public Hearing)	Family Asst Ctr., New Bedford, MA	11/8/2000
	Social Impact Informational Meeting (Public Hearing)	Narragansett Town Hall, RI	11/9/2000
Social Impact Informational Meeting (Public Hearing)	Casco Bay Terminal, Portland, ME	11/13/2000	
Social Impact Informational Meeting (Public Hearing)	Ramada Inn, Riverhead, NY	11/21/2000	
Social Impact Informational Meeting (Public Hearing)	Holiday Inn, Ellsworth, ME	12/7/2000	
Social Impact Informational Meeting (Public Hearing)	VFW, Scituate, MA	12/7/2000	

	Meeting Type	Meeting Location	Meeting Dates
2001	Council Meeting	Sheraton Ferncroft, Danvers	1/23 - 11/25/2001
	Council Meeting	Radisson, New London, CT	3/14 & 3/15/2001
	Council Meeting	Holiday Inn, Peabody, MA	5/2 & 5/3/2001
	Council Meeting	Providence Biltmore, RI	6/13 & 6/14/2001
	Oversight Committee	Sheraton Ferncroft, Danvers	1/17/2001
	Oversight Committee	Holiday Inn, Peabody, MA	2/6/2001
	Oversight Committee	Holiday Inn, Peabody, MA	2/27 & 2/28/2001
	Oversight Committee	DoubleTree Hotel, Portland, ME	6/4/2001
	Oversight Committee/Advisory Panel	Sheraton Ferncroft, Danvers	4/16/2001
	Advisory Panel	Holiday Inn, Peabody, MA	2/22/2001
	Groundfish and Capacity – Oversight Committee/Advisory Panel/PDT	Providence Biltmore, RI	6/12/2001
	PDT	NEFSC, Woods Hole, MA	1/11/2001
	PDT	Rossi's, Newburyport, MA	2/15/2001
	PDT	NEFMC, Newburyport, MA	2/21/2001
	PDT	DMF, Pocasset, MA	3/8/2001
	PDT	NEFSC, Woods Hole, MA	4/10/2001
	PDT	John Carver Inn, Plymouth, MA	5/8/2001
	PDT	DMF, Pocasset, MA	5/24/2001
	PDT/MMC	Rossi's, Newburyport, MA	8/28/2001
	PDT	NEFMC, Newburyport, MA	11/13/2001
PDT	NEFSC, Woods Hole, MA	12/3/2001	
2002	Council Meeting	Marriott Courtyard, Portsmouth NH	1/15-1/17/2002
	Council Meeting	Hilton, Mystic CT	3/19-3/20/2002
	Council Meeting	Sheraton Ferncroft Hotel, Danvers MA	5/16/2002
	Council Meeting	Samoset Resort, Rockport ME	6/24-6/26/2002
	Council Meeting	Providence Biltmore, Prov. RI	9/10-9/12/2002
	Council Meeting	Gloucester High School, MA	11/5-11/7
	Oversight Committee	Sheraton Colonial, Wakefield, MA	1/25/2002
	Oversight Committee	Holiday Inn, Mansfield, MA	2/22/2002
	Oversight Committee	Holiday Inn, Peabody, MA	3/27/2002
	Oversight Committee	Radisson Hotel, New London, CT	4/9/2002
	Oversight Committee	Holiday Inn, Mansfield, MA	4/23 & 4/24/2002
	Oversight Committee	Holiday Inn, Peabody, MA	4/30/2002
	Oversight Committee	DoubleTree Hotel, Portland, ME	5/7 & 5/8/2002
	Oversight Committee	Sheraton Colonial, Wakefield, MA	7/8/2002
	Oversight Committee	Holiday Inn, Portsmouth, NH	10/30/2002
	Oversight Committee	Holiday Inn, Mansfield, MA	12/2/2002
	Advisory Panel	Sheraton Ferncroft, Danvers, MA	5/31/2002
	Recreational Advisory Panel	Holiday Inn, Peabody, MA	4/26/2002
	Recreational Advisory Panel	Holiday Inn, Peabody, MA	6/18/2002
	PDT	NEFSC, Woods Hole, MA	2/7/2002
	PDT	NEFMC, Newburyport, MA	3/7/2002
	PDT	NEFMC, Newburyport, MA	4/15/2002
	PDT	NEFSC, Woods Hole, MA	5/29/2002
	PDT	Rossi's, Newburyport, MA	6/18/2002
	PDT	NEFSC, Woods Hole, MA	7/10/2002
PDT	NEFMC, Newburyport, MA	9/18/2002	
PDT	NEFSC, Woods Hole, MA	10/16/2002	
	Council Meeting	Courtyard by Marriott, Portsmouth, NH	1/28-1/30/2003
2003	Council Meeting	Providence Biltmore, Providence, RI	3/4-3/6/2003
	Council Meeting	Sheraton Ferncroft Hotel, Danvers, MA	4/15/2003
	Council Meeting	Tavern on the Harbor, Gloucester, MA	5/20-5/22/2003

	Meeting Type	Meeting Location	Meeting Dates
	Council Meeting	Holiday Inn, Portsmouth, NH	7/15-7/17/2003
	Oversight Committee	Holiday Inn, Mansfield, MA	1/22/2003
	Oversight Committee	Providence Biltmore, Providence, RI	3/3/2003
	Oversight Committee	Sheraton Colonial, Wakefield, MA	4/8/2003
	Oversight Committee	Holiday Inn, Peabody, MA	7/9-7/10/2003
	Advisory Panel	Holiday Inn, Peabody, MA	7/8/2003
	PDT	NEFMC, Newburyport, MA	1/17/2003
	PDT	NEFSC, Woods Hole, MA	2/25/2003
	PDT	Holiday Inn, Mansfield, MA	3/20/2003
	PDT	Holiday Inn, Mansfield, MA	4/23/2003
	Council	Tavern on the Harbor, Gloucester, MA	5/20-5/22/2003
	Council	Holiday Inn, Portland, ME	7/15-7/17/2003
	Public Hearing	Tom's River, NH	9/09/2003
	Public Hearing	Riverhead, NY	9/10/2003
	Public Hearing	South Kingstown, Ri	9/11/2003
	Public Hearing	Hyannis, MA	9/15/2003
	Public Hearing	Gloucester, MA	9/22/2003
	Public Hearing	Portsmouth, NH	9/23/2003
	Public Hearing	Ellsworth, ME	9/24/2003
	Public Hearing	Portland, ME	9/25/2003
	Public Hearing	Fairhaven, MA	9/30/2003
	Council	Holiday Inn, Peabody MA	10/21/2003
	Council	Holiday Inn, Peabody MA	11/4 – 11/6/2003

8.5 Index

- Affected Human Environment, 1, 708, 855, 860, 861, 869, 872, 882, 957, 1018, 1334, 1364
- Biological Impacts, x, xiii, 62, 64, 177, 227, 228, 272, 273, 278, 283, 287, 290, 291, 295, 296, 297, 300, 301, 303, 309, 310, 312, 313, 314, 322, 326, 334, 337, 343, 347, 349, 354, 359, 364, 365, 366, 367, 369, 372, 373, 388, 957
- Bycatch, vii, xi, xv, lxvi, 60, 121, 122, 273, 283, 369, 370, 376, 377, 379, 380, 382, 383, 393, 396, 425, 431, 448, 482, 788, 972, 988, 999, 1012, 1094, 1407
- Cod
- GB, vii, x, xv, xvii, xix, 4, 5, 32, 34, 35, 36, 37, 39, 47, 54, 55, 56, 65, 67, 76, 78, 83, 86, 114, 119, 120, 122, 134, 146, 149, 153, 155, 159, 162, 164, 165, 173, 180, 198, 200, 202, 227, 229, 234, 241, 243, 245, 246, 250, 253, 264, 284, 290, 291, 295, 300, 303, 317, 326, 333, 334, 336, 337, 342, 345, 346, 347, 348, 356, 357, 359, 360, 361, 382, 405, 406, 407, 409, 411, 413, 416, 417, 418, 419, 420, 421, 439, 441, 448, 476, 477, 568, 641, 650, 683, 786, 787, 803, 804, 839, 840, 846, 886, 887, 905, 907, 908, 909, 992, 1014, 1130, 1378, 1387, 1400, 1401, 1410, 1438
- GOM, xix, 3, 4, 5, 32, 34, 36, 42, 47, 74, 86, 91, 135, 146, 149, 155, 159, 161, 162, 164, 165, 198, 200, 224, 226, 227, 229, 235, 242, 244, 246, 251, 254, 260, 264, 300, 304, 310, 316, 317, 318, 319, 322, 323, 326, 330, 345, 346, 347, 348, 356, 357, 359, 362, 363, 401, 403, 405, 406, 407, 408, 410, 412, 413, 416, 417, 419, 420, 421, 439, 473, 475, 478, 611, 614, 624, 627, 628, 632, 638, 641, 645, 677, 767, 772, 846, 886, 887, 905, 907, 908, 964, 965, 969, 970, 1015, 1017, 1034, 1378, 1379, 1380, 1381, 1382, 1383, 1387, 1399, 1400, 1413, 1414, 1437, 1438, 1473, 1474
- Cumulative Impacts, xiv, 10, 95, 453, 858, 957, 972, 973, 974, 975, 976, 977, 978, 980, 981, 982, 983, 985, 998, 999, 1000, 1001, 1002, 1003, 1004, 1005, 1006, 1007, 1008, 1024, 1025, 1182, 1226, 1231, 1324, 1334
- Days@at@sea (DAS), 274, 278, 999, 1000, 1001, 1004, 1005, 1094
- Days-at-sea (DAS), v, vii, viii, ix, xi, xiii, xv, xvi, xvii, xviii, xix, lxvii, 3, 5, 8, 37, 53, 54, 58, 59, 60, 62, 64, 65, 66, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 79, 80, 81, 83, 84, 85, 86, 88, 89, 93, 100, 117, 118, 119, 121, 122, 125, 126, 127, 128, 129, 130, 131, 132, 134, 135, 137, 139, 140, 141, 142, 143, 144, 145, 146, 147, 149, 150, 152, 153, 154, 155, 156, 158, 159, 161, 162, 163, 165, 166, 172, 173, 174, 175, 177, 181, 197, 198, 199, 202, 203, 223, 272, 274, 278, 283, 284, 286, 287, 291, 297, 298, 299, 300, 301, 302, 303, 304, 305, 310, 311, 312, 313, 314, 315, 330, 331, 332, 333, 334, 335, 337, 338, 339, 340, 341, 342, 343, 345, 346, 349, 350, 351, 352, 354, 355, 356, 357, 381, 383, 398, 400, 401, 402, 403, 404, 405, 407, 413, 418, 427, 428, 430, 431, 432, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 446, 448, 449, 454, 455, 456, 457, 458, 471, 472, 473, 474, 475, 476, 478, 480, 481, 483, 484, 486, 488, 489, 491, 492, 558, 562, 567, 607, 608, 609, 610, 611, 614, 624, 625, 628, 630, 636, 637, 654, 655, 676, 677, 680, 683, 693, 694, 697, 698, 699, 700, 701, 702, 703, 704, 705, 706, 717, 722, 728, 730, 731, 787, 789, 790, 791, 792, 793, 794, 795, 797, 798, 799, 801, 804, 805, 808, 809, 810, 811, 812, 813, 814, 815, 816, 817, 818, 819, 820, 821, 822, 823, 825, 828, 833, 835, 836, 839, 840, 841, 842, 844, 845, 846, 847, 848, 849, 856, 857, 858, 859, 860, 861, 862, 863, 864, 865, 866, 867, 868, 871, 872, 873, 874, 875, 877, 878, 879, 880, 881, 882, 883, 889, 890, 891, 892, 893, 894, 895, 896, 897, 898, 899, 900, 901, 902, 903, 904, 905, 906, 907, 908, 909, 910, 951, 952, 955, 956, 963, 964, 965, 966, 968, 969, 970, 971, 972, 973, 974, 975, 976, 977, 979, 988, 989, 990, 993, 994, 996, 1011, 1015, 1016, 1017, 1018, 1023, 1031, 1033, 1035, 1036, 1037, 1039, 1041, 1044, 1045, 1048, 1049, 1050, 1094, 1158, 1337, 1338, 1341, 1342, 1343, 1344, 1345, 1346, 1347, 1348, 1349, 1350, 1351, 1352, 1355, 1356, 1361, 1362, 1363, 1364, 1367, 1368, 1370, 1371, 1375, 1376, 1377, 1378, 1384, 1385, 1386, 1387, 1394, 1395, 1397, 1407, 1433, 1506, 1507, 1529, 1572, 1584, 1590, 1600, 1622
- Economic impacts, xii, xiii, 8, 15, 34, 52, 53, 62, 74, 85, 125, 149, 182, 197, 225, 226, 227, 298, 300, 323, 342, 351, 354, 364, 535, 540, 543, 548, 550, 551, 553, 564, 568, 597, 607, 608, 609, 610, 611, 624, 629, 637, 655, 659, 665, 709, 712, 713, 717, 729, 735, 751, 755, 763, 767, 768, 772, 773, 778, 779, 785, 786, 787,

790, 795, 796, 803, 829, 833, 854, 856, 878, 882, 883, 888, 889, 909, 910, 911, 947, 965, 983, 984, 1016, 1019, 1023, 1025, 1026, 1032, 1033, 1038, 1334, 1440

Essential Fish Habitat, v, vii, x, xi, xiv, lxvi, lxvii, 1, 3, 6, 7, 9, 10, 11, 13, 14, 54, 61, 62, 63, 86, 92, 93, 94, 97, 98, 182, 187, 193, 204, 209, 215, 216, 225, 313, 373, 422, 443, 444, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 458, 461, 467, 470, 471, 472, 473, 477, 479, 481, 482, 483, 484, 485, 486, 488, 490, 491, 492, 493, 500, 502, 503, 504, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 523, 524, 525, 526, 527, 530, 534, 537, 538, 539, 540, 541, 542, 544, 545, 546, 547, 548, 549, 550, 551, 552, 553, 555, 556, 557, 558, 559, 561, 772, 775, 797, 834, 851, 911, 912, 944, 945, 946, 947, 948, 949, 950, 952, 953, 957, 958, 959, 963, 965, 966, 969, 973, 976, 979, 980, 981, 982, 983, 984, 1007, 1009, 1012, 1013, 1014, 1016, 1018, 1019, 1020, 1021, 1022, 1025, 1027, 1062, 1080, 1081, 1092, 1095, 1099, 1102, 1103, 1117, 1130, 1159, 1182, 1183, 1201, 1247, 1248, 1249, 1250, 1251, 1252, 1253, 1254, 1255, 1256, 1257, 1258, 1259, 1260, 1261, 1262, 1263, 1264, 1265, 1266, 1267, 1268, 1269, 1270, 1271, 1272, 1273, 1274, 1275, 1276, 1277, 1278, 1279, 1280, 1281, 1282, 1283, 1284, 1285, 1286, 1287, 1288, 1289, 1290, 1291, 1292, 1293, 1294, 1295, 1296, 1297, 1298, 1299, 1300, 1301, 1302, 1303, 1304, 1305, 1306, 1307, 1308, 1309, 1310, 1312, 1313, 1314, 1315, 1316, 1317, 1318, 1319, 1324, 1325, 1326, 1327, 1329, 1330, 1332, 1333, 1357

Exclusive Economic Zone, xvi, lxvii, 72, 128, 773, 841, 1030, 1095, 1148, 1183, 1187, 1188, 1189, 1202, 1285, 1286, 1326, 1327, 1439

Habitat impacts, 62, 64, 77, 374, 446, 447, 448, 449, 455, 456, 457, 461, 470, 472, 480, 481, 482, 483, 484, 485, 486, 487, 491, 548, 555, 1014, 1319, 1324

Habitat Impacts, xi, 446, 458, 472, 476, 479, 480, 481, 487, 488, 489, 957

Haddock
 GB, 3, 5, 32, 34, 36, 42, 47, 69, 164, 177, 230, 233, 235, 242, 244, 247, 251, 255, 261, 265, 290, 291, 300, 325, 333, 344, 355, 419, 483, 786, 790, 804, 883, 1380, 1386, 1387, 1415
 GOM, 32, 36, 37, 42, 47, 164, 177, 233, 310, 320, 321, 326, 344, 355, 419, 884, 1136, 1386, 1410

Haddock: GB, 1003

Halibut, 21, 25, 33, 38, 49, 51, 102, 107, 110, 116, 180, 331, 332, 338, 339, 340, 341, 349, 350, 353, 358, 451, 477, 502, 511, 556, 1135, 1184, 1185, 1253, 1307, 1312, 1317, 1409

Magnuson-Stevens Fishery Conservation and Management Act, iii, v, lxvii, 1, 3, 4, 6, 7, 8, 10, 11, 13, 15, 19, 20, 31, 32, 34, 35, 36, 51, 59, 64, 69, 91, 103, 104, 105, 108, 114, 145, 182, 197, 204, 228, 272, 273, 291, 312, 333, 334, 372, 379, 380, 473, 491, 511, 546, 574, 787, 854, 858, 1011, 1016, 1020, 1022, 1025, 1026, 1027, 1030, 1133, 1182, 1247, 1326, 1330, 1334, 1384, 1407, 1436

Marine mammals, 380, 426, 436, 440, 960, 965, 1025, 1088, 1091, 1195

Mid-Atlantic, 1030

Myxine glutinosa an., 1027, 1028

National Environmental Policy Act, 1024, 1025

Ocean pout, 42, 47, 49, 502, 1165, 1265

Plaice, American, v, xix, 21, 24, 32, 33, 37, 38, 43, 47, 49, 50, 51, 83, 87, 88, 102, 106, 109, 114, 116, 147, 159, 163, 180, 231, 237, 241, 248, 252, 256, 259, 262, 266, 314, 315, 320, 323, 324, 325, 327, 329, 331, 332, 334, 335, 338, 339, 340, 341, 343, 349, 350, 351, 353, 358, 371, 383, 423, 451, 471, 476, 477, 502, 511, 513, 515, 556, 565, 967, 1053, 1054, 1055, 1056, 1057, 1058, 1099, 1101, 1129, 1135, 1160, 1164, 1165, 1184, 1185, 1250, 1307, 1312, 1317, 1325, 1389, 1409, 1420, 1421, 1431, 1432, 1435, 1502, 1511, 1519, 1521, 1523, 1530, 1532, 1534, 1536, 1539, 1542, 1550, 1553, 1555, 1562, 1564, 1573, 1576, 1582, 1584, 1586, 1588, 1592, 1594, 1596, 1602, 1604, 1607, 1610, 1618, 1624, 1630, 1633, 1639, 1644, 1647, 1652, 1655, 1659

Pollock, xix, 21, 25, 33, 38, 49, 50, 87, 102, 107, 110, 116, 147, 159, 163, 164, 177, 180, 234, 270, 314, 315, 327, 329, 331, 332, 335, 338, 339, 340, 341, 343, 344, 349, 350, 351, 353, 355, 358, 371, 377, 419, 451, 476, 477, 502, 512, 556, 802, 1135, 1165, 1184, 1185, 1267, 1308, 1313, 1317, 1335, 1409, 1435

Practicability, 98, 537, 538, 539, 546, 548

Protected species, 431

Redfish, xix, 21, 25, 33, 38, 42, 47, 49, 50, 87, 102, 107, 110, 116, 147, 159, 163, 180, 232, 238, 249, 259, 263, 267, 314, 315, 324, 325, 327, 329, 331, 332, 335, 338, 339, 340, 341, 343, 349, 350, 351, 353, 358, 371, 451, 476,

477, 502, 510, 512, 513, 515, 518, 557, 1135,
1184, 1185, 1270, 1308, 1313, 1317, 1409
Social Impact Analysis, lxviii, 854, 855, 860,
870, 884, 887, 888, 907, 908, 1493
Social impacts, xiii, 225, 857, 861, 881, 910, 957
Special Access Program, vii, x, xii, xv, xix, lxviii,
54, 58, 61, 62, 63, 64, 66, 68, 73, 80, 85, 122,
273, 274, 290, 295, 382, 383, 398, 399, 432,
448, 483, 610, 789, 790, 839, 889, 973, 988,
1036, 1037, 1044
Status Determination Criteria, vi, 15, 18, 19, 22,
27, 101, 103, 105, 446, 482
White hake, 37, 47, 477, 503, 1272, 1412
Windowpane flounder, 42, 47, 1274
Winter flounder
GB, 164, 177, 284, 290, 292, 333, 344, 349,
355, 419, 964, 1386, 1387, 1410
GOM, 23, 164, 177, 239, 270, 271, 326, 344,
355, 403, 419, 1011, 1133, 1413, 1427,
1428, 1429

SNE/MA, 32, 42, 47, 66, 114, 232, 237, 241,
243, 249, 253, 257, 259, 263, 267, 287, 333,
349, 359, 884, 973, 1011, 1133, 1386, 1413,
1418, 1419
Witch flounder, xix, 87, 147, 159, 163, 269, 325,
334, 476, 477, 503, 1165, 1276, 1386, 1422,
1423
Yellowtail flounder
CC/GOM, x, xix, 32, 35, 36, 37, 39, 47, 48, 83,
86, 156, 159, 162, 200, 230, 236, 241, 243,
245, 248, 250, 253, 256, 259, 262, 266, 300,
325, 327, 329, 334, 336, 348, 381, 403, 405,
406, 407, 409, 412, 414, 439, 834, 886, 905,
1011, 1130, 1413, 1424, 1425
SNE/MA, xix, 21, 32, 37, 47, 48, 83, 86, 102,
156, 159, 162, 164, 231, 236, 243, 245, 247,
252, 255, 259, 261, 265, 300, 325, 327, 329,
331, 332, 338, 339, 340, 341, 349, 350, 353,
356, 357, 358, 405, 406, 407, 409, 412, 415,
418, 419, 420, 421, 439, 883, 886, 905,
1011, 1130, 1133, 1135, 1136, 1413, 1426

INTENTIONALLY BLANK

INTENTIONALLY BLANK

INTENTIONALLY BLANK